

Review Article

The Effects that the Current Climate Crisis have on the Biogeography and Environment, Needed Adaptations and Conservation

Ida Krogsgaard Svendsen

Centre for Ecological and Evolutionary Synthesis (CEES), Department of Biosciences, University of Oslo, Oslo, Norway

Email address:

Idsve10@gmail.com

To cite this article:

Ida Krogsgaard Svendsen. The Effects that the Current Climate Crisis have on the Biogeography and Environment, Needed Adaptations and Conservation. *American Journal of BioScience*. Vol. 8, No. 1, 2020, pp. 20-27. doi: 10.11648/j.ajbio.20200801.14

Received: March 7, 2020; **Accepted:** March 23, 2020; **Published:** April 13, 2020

Abstract: This study is a literature review aiming to give a summary of the effects that the current anthropogenic caused climate crisis has on the biogeography and environment, and further give examples of likely future adaptations and needed conservation work. This study is based on scientific articles, primary from Web of Science and Google Scholar. The biodiversity is under pressure due to climate changes, the average species extinction is currently two to three orders of magnitude higher than the normal background extinction, and faster than the rate of origination. This development follows the predictions of The Red Queen Hypothesis that every species must constantly evolve due to environmental changes in order to avoid extinction. The natural environments are changing due to e.g. increased extreme weather events and ocean acidification. The increased heating is causing drought, and adaptations of the biota is needed, like more drought resistant flora and fauna with the ability to undergo estivation. The increased oceanic acidity can cause the shells of calcifying organisms to dissolve. These organisms will need to either spend energy on increased calcification or develop in a way so they can carry out live with lesser calcification. If organisms cannot develop, they are likely to migrate to colder regions. In the ocean this means towards polar areas and to greater depths, and in the terrestrial environment it is pole wards and to greater altitudes. Conservation is needed, and there are multiple options. Ex situ might be the only option for species whose natural habitat will be forever gone if the development of the climate change continues as present. To carry out conservation to infinity is unrealistic, and we are at a point where climate change is threatening our food security. It is possible to both slow down the current climate crisis and counteract its consequences.

Keywords: Climate Change, Biogeography, Conservation, Extreme Weather, Adaptations, Migration

1. Introduction

This paper provides the current knowledge about how the current anthropogenic climate crises is affecting the biogeography and our environment. Subjects that this paper is touching is: (1) the needed adaptations that species will need to undergo in order to avoid extinction, (2) ocean acidification and its consequences, for especially calcifying organisms, (3) problems with more extreme weather, like desertification in some areas while heavier rain will be more frequent in other, (4) the needed migration pole wards, to greater altitudes and greater oceanic depth in order to stay in the preferred thermal conditions, and (5) our food security is threatened by a climate changed induced mismatch between pollination and flowering

season.

2. A Summary of the Current Climate Change Crises and Needed Adaptations

2.1. Extinction and the Need to Evolve

Extinction is a common phenomenon, and as described by The Red Queen Hypothesis, species will constantly have to respond to changing environmental conditions in order to avoid going extinct [1, 2]. The biodiversity on Earth have been shaped by the climatic and environmental conditions

throughout evolutionary time [3], and organisms that cannot cope with the changing environmental conditions will need to either adjust or migrate to more thermal suitable habitats [4]. To survive, species need to constantly evolve fast enough to be suitable adapted to the environment, and have a fitness that can compete with predators, parasites etc., who also are continually evolving [5, 6, 7]. Meaning that for species to exist, they must minimally be well-designed, and if better designed they might be more suited for future circumstances [6], maladapted species will be eliminated by natural selection [5].

2.2. Be Adapted to Avoid Extinction

Being adapted to the environment in order to avoid extinction is essential, both with respect to the present climate changes due to anthropogenic activities and in an historical context. It is estimated that 95-99% of the four billion species that have evolved on- and inhabited the Earth over the last 3.5 billion years have suffered from extinction [8-10]. Extinction has always occurred, but currently the average species extinction is 0.72% per year, which is two to three orders of magnitude higher than the normal background extinction, and faster than the rate of origination [11, 9, 12].

The big driver for this extinction is climate changes, which is currently more rapid than what have ever occurred during recorded history [1, 13]. The expected global mean surface temperature for 2100 will be between 3.7°C and 4.8°C above the average for the period 1850-1900 if no drastic effort is taken to reduce the emission of greenhouse gases [14].

Since the industrial revolution, which started in the 19th century, the level of atmospheric carbon dioxide has increased from about 281 ppm to more than 400 ppm, which is the highest in recorded history [15, 16]. There are multiple greenhouse gasses contributing to climate change but the release of carbon dioxide to the atmosphere is especially critical because the worldwide output is so enormous [17, 18].

3. Ocean Acidification and Its Consequences

3.1. Oceanic Sink and Problems for Calcifying Species

The ocean is a major sink for atmospheric carbon dioxide [19], and accounts for 48% of the uptake [15], but someday the oceans capacity for taking up carbon dioxide might be reached, meaning that a further uptake will not be possible [15, 20]. Ocean acidification will expose many species to conditions that they have not been exposed to during their evolutionary history, and for this reason is not adapted to, which might cause extinction [20].

Carbon dioxide can enter some organisms and changes the pH of their body fluids. When organisms use energy on regulating pH, it can have consequences for growth rate and reproductive capacity [21-24]. The effect of ocean acidification is severe on calcifying organisms, because it can decrease their carbonate ion concentration to levels where

shells dissolve [25, 26]. This have consequences for many ecosystems since reef-building corals and calcified pteropods are affected. Pteropods have essential roles in food webs of various marine ecosystems [27]. The coral reefs provide breeding grounds for multiple fish, protect sea grasses from wave stress, and protect shorelines from erosion and flooding [28, 20, 29].

3.2. Problems with the Stress from Heat

The heat is also a problem for corals, because they cannot make their own photosynthesis, they are dependent on their photosynthetic algal endosymbionts (*zooxanthellae*) in order to survive. But the thermal tolerance of *zooxanthellae* has been exceeded in many reefs, which have led to widespread coral bleaching [17, 30]. If the carbon dioxide emission continues as present, 70% of all known tropical reef locations will be gone at the end of this century [20].

3.3. Possible Adaptation the Acidic Environment

A species which have successfully undergone evolutionary adaptations to live in an acidified aquatic environment is *paedocypris* (*Paedocypris progenetica*), which is a miniature fish from Southeast Asia. They are native to caves with a naturally highly acidic environment and is adapted to these environments in the way that they are capable to live and function with a poorly mineralized skeleton [31].

4. Problems with Extreme Weather

4.1. Desertification

The global warming is causing an increase in the frequency, duration, and severity of drought in some regions [32]. A Drought climate will often result in increased wildfires, and thus vegetation will be lost [33-35]. Since plants remove atmospheric carbon dioxide, the terrestrial carbon sink will decrease due to increased wildfire [18, 36]. It is estimated that the global terrestrial carbon sink absorbs a range of 2.0 to 3.4 Pg. carbon per year [36]. If these pools of carbon become released back into the atmosphere it will contribute to acceleration of further global warming [32].

4.2. Not a New Phenomenon, and Some Species Are Already Adapted

Drought and wildfires are not new phenomena and has influenced the atmosphere and biosphere for millions of years [37]. In climate zone ecosystems like the Mediterranean-type, the native flora is particularly adapted to drought. These habitats are dominated by woody species, and plants that have a large amount of underground reserves to re-sprout and soil seed-banks [38, 39, 34, 40]. Lungfish, many insects, and desert snails, can escape periods with drought by using estivation, which is a state where the metabolic rate is substantially reduced and the body is at rest, despite the high environmental temperatures [41, 42]. They can survive estivation without any excretion of nitrogenous

waste products [42]. The abilities for osmoregulation seem to be adaptive within a species at the population level, and there have been shown significant anatomical differences in the relative medullary thickness between kangaroo rats living under drought conditions, compared with more moisture conditions [43]. Species of mammals that live under constant drought stress have long loop of Henle in proportion to their body size, which makes them able to produce hyperosmotic urine. This is a favorable adaptation in where they lose very little water when excreting excess salts [42].

4.3. The Extreme Weather also Comes with the Problems of Flooding

The global warming is also associated with increased flooding in warmer maritime regions because the higher temperatures is increasing the evaporation from water surfaces [44, 45, 35]. The hazards that follows are likely to increase in the future with the predicted climate change [46]. The reason why the flooding increases in some areas, while drought is a severe problem in other, is because areas with closer proximity to aquatic areas will experience more rainfalls caused by evaporation, which will not happen in regions distant to aquatic areas [47, 48].

5. Migration

5.1. Some Species Need to Follow Their Thermocline Gradient

Many taxonomic groups worldwide have migrated pole ward and upslope to escape the heat [49, 50]. When migrating to high-altitude areas the adaptations to hypoxia is the main physiological challenge that threatens survival [51], and animals adapted to high altitudes in general have hemoglobin with a higher oxygen affinity, and larger heart and lungs [52, 53].

5.2. In the Ocean There Are Migration to Greater Depths

Due to increased ocean temperature multiple marine species are migrating pole wards or to greater depths [54, 55], and a globally large-scale invasion into colder marine waters is expected in the future [56]. This migration might cause conflicts between migrants and native species in receiving areas [57].

6. Pollination

6.1. Climate Change Is Threatening Our Food Security

The interactions between plants and pollinators are among the most ecologically important mutualistic relationships in nature, and fundamental for the global biodiversity, and further an important selective force during evolution [5-60]. But unfortunately, global warming is disturbing this relationship [61, 60].

6.2. Climate Change Induced Mismatches Gives Problems for the Pollination

For multiple species of plants there is a strong and complex relation between air temperature and pollen release [62]. The global warming leads to an earlier onset of the pollen season in some species [62], which can result in temporal mismatches between plant and pollinator [63-65]. In some places, like Arctic Greenland, this have led to a decline in the overlap between the pollen transfer season and the flowering season [60]. Shifted flowering seasons have made some pollinators incapable to track all their ancestral hosts flowers [63, 66]. These mismatches contribute to decreased pollination efficiency and have led to measurable impacts on the plant's reproductive success [67, 64, 60]. In some cases, the mismatches lead to extinctions of both plant and pollinator [68]. It is likely that specialized pollinators will end up with no food plants, while pollinators that are more generalist will just experience diet reductions following phenological shifts [63]. Animals pollination is fundamental for the food security, because one-third of agricultural production depends on it [69]. A decline will lead to an increased demand for agricultural land [70]. Further is the pollination also essential for species that feeds on flowering plant [71, 72].

7. Conservation and Sustainability

7.1. Conservation Is a Huge Task

Conservatory work has through times resulted in both victories and failure, and currently the rapid climate change is placing the future of the conservation legacy in a vulnerable position [73, 74]. Conservation is currently a huge task because the distribution of many species and ecological processes are changing much faster than in the past [75]. The rate of climate change is likely to be so fast that the genetic adaptation of many species will not be able to cope [76, 77], nor is it likely for many species that their migration will be fast enough [78]. Unfortunately, along with that every endemic species is irreplaceable, the funding for conservation action is limited while the resources needed is increasing due to climate change [79, 80, 76, 81].

7.2. Consequences of Changed Ecosystems due to Anthropogenic Activities

Over the past five decades the anthropogenic activities have changed multiple ecosystems more rapidly than have ever before been recorded, which have caused irreversible loss of biodiversity [82]. Deforestation and agricultural expansion have resulted in destruction and fragmentation of many habitats, which have resulted in endangerment of many species [83, 84]. This fragmentation causes island conditions where many of the inhabiting species have smaller population size, and are geological limited in their options for dispersal, which is likely to cause extinctions [85, 80, 86]. Mainly due to habitat fragmentations 80% of vertebrate taxa have become globally endangered [87], and between 1 and 2 million km² of

tropical forests are destroyed by anthropogenic actions every decade [82, 84]. Destruction of the tropical forest is detrimental because it stores around 46% of the world's living terrestrial carbon pool, and the tropical soils about 11.55% [88, 89]. If the land-use change and deforestation activities were limited, the global greenhouse gas emission and pressure on the biodiversity will be reduced [18, 90].

7.3. Sustainability Is Likely to Be the Solution

Decreasing the greenhouse gas emission through a more sustainable anthropogenic consumption and way of living is possible. A more plant-based diet is more sustainable than a meat-based diet, partly because meat-production consumes a large portion of the crops and increases the problem of water shortages. It takes around 1,000 tons of water to produce 1 ton of grain, and it can take up to 8 kg of grain to produce just 1, 2, or 4 kg of beef, poultry or pork, respectively [91-93]. Resources like paper and fibers can be recycled [94], and the use of solar energy decreases the use of other energy sources [95-97].

7.3.1. The Need to Relocate Species and Establish Protected Areas

By establishing protected areas, the habitat will be maintained, which is a sufficient method to prevent losing habitat while protecting endangered species [79, 84, 98, 11]. Under rapidly changing climatic conditions the challenge of maintaining and restoring habitats can be very difficult [99, 100], and conservation workers must often use more proactive methods, such as removing invasive competitors or relocating populations to new habitat [79]. Relocation of species to areas outside their present or historical range, where their survival is expected to be favored, is a tool that can be used to conserve biodiversity under the future climatic changes. By using this method, the problem of dispersal barriers is solved, meaning that it is letting species survive climate change by altering their geographic distributions [101-104]. This method can also be used to relocate some species of turtles into cooler regions in order to counteract the climate change induced female-biased sex ratio of their offspring [105]. Relocation of species is not exclusively success, relocated species are in general likely to have a higher mortality rate than what is found naturally in the wild [106].

7.3.2. Ex Situ Conservation Might Be the only Solution for Some Species

Ex situ conservation is another important tool, which basically is when species are moved into artificial ecosystems like zoos, gardens, or aquariums. This way of conservation is less risky than relocation of species in the nature. Ex situ conservation might be the only option for saving species that live near the geographic end of climatic gradients, like polar and alpine species [107].

8. Discussion and Conclusion

The current climate crisis is caused by anthropogenic activities and have led to disruptions of multiple ecosystems.

Among greenhouse gases the main driver is the emissions of carbon dioxide [22]. The atmospheric carbon dioxide can be absorbed by carbon sinks, like the ocean and vegetations. Due to deforestation and urbanization, the terrestrial carbon sink is decreasing, and the capacity for oceanic uptake might reach its limit [15, 20, 32]. For these reasons actions like reforestation and limiting the output of atmospheric carbon dioxide is of importance.

Species might face extinction if they cannot develop fast enough to cope with the changing environment [7]. What differs between historical mass-extinctions and the current, is that the current is anthropogenic caused. Currently the average species extinction is 0.72% per year, which is two to three orders of magnitude higher than the normal background extinction [12], since this is faster than the rate of origination the biodiversity on Earth is declining.

Due to the acidification of the oceans many marine organisms will be exposed to conditions that they are not adapted to, which might be detrimental for their existence [20]. Under acidification some organisms reduce their metabolic rate, which have the drawback of reduced reproduction [21]. Especially calcifying organisms will need adaptations to a more acidic environment, either they will have to increase their rate of calcification to compensate for the acidification or they will have to manage life in a way similar to *Paedocypris progenetica*.

Many terrestrial ecosystems will become more drought [32]. Adaptations that likely will be more common in the future as a result of drought includes a flora dominated by woody species, and plants that have a large amount of underground reserves to re-sprout and soil seed-banks [38], [39]. Species like lungfish can escape periods with drought by using estivation, and many desert animals have evolved long loop of Henle in proportion to their body size [42]. These adaptations will likely be more common in the future due to the increased heat stress and more frequent occurrence of extreme weather events.

Some species migrate to escape the heating. Oceanic species can migrate pole wards or to greater depths [54, 55]. While terrestrial species can migrate pole wards and to greater altitudes [49, 50]. When migrating to greater heights, adaptations such as hemoglobin with a higher oxygen affinity, larger heart, and lungs are needed [52, 53]. It might be questionable if all the needed adaptations for migrating to greater heights can occur as fast as needed, and also when new species invades an already occupied area in the heights it will undoubtedly lead to conflicts between established and invading species.

The climatic changes have also disturbed the pollination. In some areas it has shifted the flowering season and made pollinators incapable to track all their ancestral hosts flowers [63, 66]. In some cases, the climate induced mismatches have led to extinctions of both plant and pollinator [68]. The decline in the function of pollination is catastrophic for the future because a big part of our food security will be threatened.

Conservation actions are needed, and there are multiple options, like establishment of protected areas, relocation of species, and ex situ conservation [101, 107, 84, 98, 108]. Relocation of species might not be possible to infinity, since at

some point it will be impossible to find a climatic suitable habitat in a changing world. In extreme cases like when the natural habitat is forever gone, e.g. polar and alpine habitats, ex situ conservation will be the only possibly option in order to avoid extinction. It is not possible to conserve to infinity, and at a point we will need to take action to change the reason why we have all the problems with global warming that leads to loss of habitats and biodiversity and even threatens our food security due to decreased efficiency of pollination. It is possible to slow down the crises and even counteract it. We can increase the carbon sink by actions like reforestation and we can decrease the anthropogenic emission by changing our way of living and consuming.

Concluding Remarks

The core problem is the emission of greenhouse gasses, but we keep trying to fix its consequences instead of stopping the problem. We need to be proactive, like stop to drill up fossil fuel and burn it to have energy, and we need to stop the continuing deforestation, we need the vegetation to suck out the carbon dioxide from the atmosphere so we can counteract the global warming. There are solutions on how to carry out life in a more sustainable way, but for it to work it requires actions. If we keep misusing the globe and its resources which we depend on the next candidate for extinction might be humans.

Acknowledgements

I want to thank my two supervisors, Nils Christian Stenseth (Professor and director of the Centre for Ecological and Evolutionary Synthesis (CEES) at the University of Oslo) and Johan Petter Dahlgren (Associate Professor of Plant Demography at the Department of Biology and the Interdisciplinary Centre on Population Dynamics (CPop), University of Southern Denmark). Their supervision during my Master's Thesis, which this article is based on, have been essential for the outcome and made me grow as a scientist. I am truly thankful.

References

- [1] Hoffmann A. A., Sgrò C. M. (2011) Climate change and evolutionary adaptation. *Nature*, 470, 479-485.
- [2] Stenseth N. C. (1979) Where have all the species gone? On the nature of extinction and the Red Queen Hypothesis. *Oikos*, Vol. 33, No. 2, pp. 196-227.
- [3] Wiens J. J. (2011) The causes of species richness patterns across space, time, and clades and the role of "ecological limits". *Q. Rev. Biol.* 86: 75-96.
- [4] Blier P. U., Lemieux H., Pichaud N. (2014) Holding our breath in our modern world: will mitochondria keep the pace with climate changes? *Can. J. Zool.* Vol. 92.
- [5] Castrodeza C. (1979) NON-PROGRESSIVE EVOLUTION, THE RED QUEEN HYPOTHESIS, AND THE BALANCE OF NATURE. *Acta Biotheoretica* 28, 1: 11-18.
- [6] McCune A. R. (1982) On the Fallacy of Constant Extinction Rates. *Evolution*, Vol. 36, No. 3, pp. 610-614.
- [7] Futuyma D. J. (2013) *evolution*, third edition. Sinauer Associates, Inc. USA.
- [8] Stanley S. M. (1987) *Extinction*. New York: Scientific American Books, Inc.
- [9] Barnosky A. D., Matzke N., Tomiya S., et. al. (2011) Has the Earth's sixth mass extinction already arrived? *Nature*, Vol 471.
- [10] Jablonski D. (2012) Paleontology in the twenty-first century. In D. Sepkoski and M. Ruse (Eds.). *The Paleobiological revolution*. Chicago University Press. pp. 471-517.
- [11] Hoffmann M., Hilton-Taylor C., Angulo A., et. al. (2010) The Impact of Conservation on the Status of the World's Vertebrates. *Science* Vol 330.
- [12] Mora C., Rollo A., Tittensor D. P. (2013) Comment on "Can We Name Earth's Species Before They Go Extinct?" *Science* Vol 341.
- [13] Santos A. M. C., Field R., Ricklefs R. E. (2016) New directions in island biogeography. *Global Ecology and Biogeography*, 25, 751-768.
- [14] IPCC (2014) *Climate Change 2014, synthesis report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, R. K. Pachauri and L. A. Meyer (eds.)]. IPCC, Geneva, Switzerland, 151.
- [15] Sabine C. L., Feely R. A., Gruber N., et. al. (2004) The Oceanic Sink for Anthropogenic CO₂. *Science*, Vol. 305.
- [16] Horwitz R., Hoogenboom M. O., Fine M. (2017) Spatial competition dynamics between reef corals under ocean acidification. *Scientific Reports*, 7: 40288.
- [17] Harley C. D. G., Hughes A. R., Hultgren K. M., et. al. (2006) The impacts of climate change in coastal marine systems. *Ecology Letters* 9, 228-241.
- [18] Malmshiemer R. W., Heffernan P., Brink S., et. al. (2008) Forest Management Solutions for Mitigating Climate Change in the United States. *Journal of Forestry*.
- [19] Watson A. J., Schuster U., Bakker D. C. E., et. al. (2009) Tracking the Variable North Atlantic Sink for Atmospheric CO₂. *Science*, Vol. 326.
- [20] Riebesell U., Körtzinger A., Oschlies A. (2009) Sensitivities of marine carbon fluxes to ocean change. *PNAS*, Vol. 106, no. 49.
- [21] Pörtner H. O., Langenbuch M., Reipschläger A. (2004) Biological Impact of Elevated Ocean CO₂ Concentrations: Lessons from Animal Physiology and Earth History. *Journal of Oceanography*, Vol. 60, pp. 705-718.
- [22] World Ocean Review. Bollmann M., Bosch T., Colijn F., Ebinghaus R., Froese R., Güssow K., Khalilian S., Krastel S., et. al. (2010) world ocean review, *Living with the ocean*. Maribus.
- [23] Busch D. S., Maher M., Thibodeau P., McElhany P. (2014) Shell Condition and Survival of Puget Sound Pteropods Are Impaired by Ocean Acidification Conditions. *PLOS ONE*, Volume 9, Issue 8, e105884.

- [24] Bednaršek N., Harvey C. J., Kaplan I. C., Feely R. A., Možina J. (2016) Pteropods on the edge: Cumulative effects of ocean acidification, warming, and deoxygenation. *Progress in Oceanography* 145, 1–24.
- [25] McNeil B. I., Matear R. J. (2008) Southern ocean acidification: a tipping point at 450-ppm atmospheric CO₂. *Proceedings of the National Academy of Sciences of the United States of America* 105 (48), 18860–18864.
- [26] Lischka S., Büdenbender J., Boxhammer T., Riebesell U. (2011) Impact of ocean acidification and elevated temperatures on early juveniles of the polar shelled pteropod *Limacina helicina*: mortality, shell degradation, and shell growth. *Biogeosciences*, 8, 919–932.
- [27] Comeau S., Gorsky G., Jeffree R., Teyssié J.-L., Gattuso, J.-P. (2009) Impact of ocean acidification on a key Arctic pelagic mollusc (*Limacina helicina*). *Biogeosciences* 6, 1877–1882.
- [28] Moberg F., Folke C. (1999) Ecological goods and services of coral reef ecosystems. *Ecol. Econ.* 29: 215–33.
- [29] van de Koppel J., van der Heide T., Altieri A. H., et. al. (2015) Long-Distance Interactions Regulate the Structure and Resilience of Coastal Ecosystems. *Annu. Rev. Mar. Sci.*, 7: 139–58.
- [30] Jones A. M., Berkelmans R., van Oppen M. J. H., Mieog J. C., Sinclair W. (2008) A community change in the algal endosymbionts of a scleractinian coral following a natural bleaching event: field evidence of acclimatization. *Proc. R. Soc. B.*, 275, 1359–1365.
- [31] Britz R., Conway K. W. (2009) Osteology of *Paedocypris*, a Miniature and Highly Developmentally Truncated Fish (Teleostei: Ostariophysi: Cyprinidae). *Journal of Morphology* 270: 389–412.
- [32] Allen C. D., Macalady A. K., Chenchouni H., Bachelet D., et. al (2010) A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management* 259, 660–684.
- [33] Grose M. R., Fox-Hughes P., Harris R. M. B., Bindoff N. L. (2014) Changes to the drivers of fire weather with a warming climate—a case study of southeast Tasmania. *Climatic Change*, 124: 255–269.
- [34] Harvey B. J. (2016) Human-caused climate change is now a key driver of forest fire activity in the western United States. *PNAS*, Vol. 11, no. 42, 11649–11650.
- [35] Garbolino E., Sanseverino-Godfrin V., Hinojos-Mendoza G. (2016) Describing and predicting of the vegetation development of Corsica due to expected climate change and its impact on forest fire risk evolution. *Safety Science* 88, 180–186.
- [36] Pan Y., Birdsey R. A., Fang J., et al. (2011) A Large and Persistent Carbon Sink in the World's Forests. *Science*, Vol 333.
- [37] Pechony O., Shindell D. T. (2010) Driving forces of global wildfires over the past millennium and the forthcoming century. *PNAS*, Vol 107, no. 45, eds. Chapin F. S., University of Alaska.
- [38] Vennetier M., Ripert C. (2009) Forest flora turnover with climate change in the Mediterranean region: A case study in Southeastern France. *Forest Ecology and Management* 258S, S56–S63.
- [39] West A. G. (2009) Climate change, drought and biodiversity: Anecophysiological perspective. *SAAB Annual Meeting Abstracts*.
- [40] Mitsopoulos I., Mallinis G., Karali A., Giannakopoulos C., Arianoutsou M. (2016) Mapping fire behaviour under changing climate in a Mediterranean landscape in Greece. *Reg Environ Change*, 16: 1929–1940.
- [41] Rasmussen O. (2005) *Håndbog i Biologiske fagtermer*, 2. edition. Gads Forlag.
- [42] Willmer P., Stone G., Johnston I. (2011) *Environmental Physiology of Animals*. Second edition, Blackwell Publishing.
- [43] Schwimmer H., Haim A. (2009) Physiological adaptations of small mammals to desert ecosystems. *Integrative Zoology*, 4: 357–366.
- [44] Whetton P. H., Fowler A. M., Haylock M. R., Pittock A. B. (1993) IMPLICATIONS OF CLIMATE CHANGE DUE TO THE ENHANCED GREENHOUSE EFFECT ON FLOODS AND DROUGHTS IN AUSTRALIA. *Climatic Change* 25: 289–317.
- [45] Hirabayashi Y., Kanae S., Emori S., Oki T., Kimoto M. (2008) Global projections of changing risks of floods and droughts in a changing climate. *Hydrological Sciences Journal*, 53: 4, 754–772.
- [46] Quataert E., Storlazzi C., van Rooijen A., Cheriton O., van Dongeren A. (2015) The influence of coral reefs and climate change on wave-driven flooding of tropical coastlines. *Geophysical Research Letters*, 42, 6407–6415.
- [47] Natugonza V., Ogotu-Ohwayo R., Musinguzi L., Olokotum M., Naigaga S., Kitabona J. (2016) Implications of climate warming for hydrology and water balance of small shallow lakes: A case of Wamala and Kawi, Uganda. *Aquatic Ecosystem Health & Management*, 19: 4, 327–335.
- [48] Mac Nally R., Horrocks G. F. B., Lada H. (2017) Anuran responses to pressures from high-amplitude drought–flood–drought sequences under climate change. *Climatic Change*, 141: 243–257.
- [49] Feeley K. J., Rehm E. M. (2012) Amazon's vulnerability to climate change heightened by deforestation and man-made dispersal barriers. *Global Change Biology*, 18, 3606–3614.
- [50] Bauera A., Farrella R., Goldblumb D. (2016) The geography of forest diversity and community changes under future climate conditions in the eastern United States. *Écoscience*, Vol. 23, NOS. 1–2, 41–53.
- [51] Brown E. A. (2012) Genetic explorations of recent human metabolic adaptations: hypotheses and evidence. *Biol. Rev.*, 87, pp. 838–855.
- [52] Monge C., León-Velarde F. (1991) Physiological Adaptation to High Altitude: Oxygen Transport in Mammals and Birds. *PHYSIOLOGICAL REVIEWS*, the American Physiological Society Vol. 71, No. 4, 0031–9333.
- [53] Zhang Q., Gou W., Wang X., et. al. (2016) Genome Resequencing Identifies Unique Adaptations of Tibetan Chickens to Hypoxia and High-Dose Ultraviolet Radiation in High-Altitude Environments. *Genome Biol. Evol.* 8 (3): 765–776.

- [54] Dulvy N. K., Rogers S. I., Jennings S., Stelzenmüller V., Dye S. R., Skjoldal H. R. (2008) Climate Change and Deepening of the North Sea Fish Assemblage: A Biotic Indicator of Warming Seas. *Journal of Applied Ecology*, Vol. 45, No. 4, pp. 1029-1039.
- [55] Kortsch S., Primicerio R., Fossheim M., Dolgov A. V., Aschan M. (2015) Climate change alters the structure of arctic marine food webs due to poleward shifts of boreal generalists. *Proc. R. Soc. B* 282: 20151546.
- [56] Holopainen R., Lehtiniemi M., Meier H. E. M., et. al. (2016) Impacts of changing climate on the non-indigenous invertebrates in the northern Baltic Sea by end of the twenty-first century. *Biol Invasions*, 18: 3015-3032.
- [57] Reuveny R. (2007) Climate change-induced migration and violent conflict. *Political Geography* 26, 656-673.
- [58] Fiz O., Vargas P., Alarcón M., Aedo C., et. al. (2008) Phylogeny and Historical Biogeography of Geraniaceae in Relation to Climate Changes and Pollination Ecology. *Systematic Botany*, 33 (2): 326-342.
- [59] Giannini T. C., Acosta A. L., Garófalo C. A., et. al. (2012) Pollination services at risk: Bee habitats will decrease owing to climate change in Brazil. *Ecological Modelling* 244 (2012) 127-131.
- [60] Schmidt N. M., Mosbacher J. B., Nielsen P. S., Rasmussen C., Høye T. T., Roslin T. (2016) An ecological function in crisis? The temporal overlap between plant flowering and pollinator function shrinks as the Arctic warms. *Ecography* 39: 1250-1252.
- [61] Bolmgren K., Eriksson O. (2015) Are mismatches the norm? Timing of flowering, fruiting, dispersal and germination and their fitness effects in *Frangula alnus* (Rhamnaceae). *Oikos* 124: 639-648.
- [62] Tedeschin E., Rodríguez-Rajo F. J., Caramiello R., Jato V., Frenguelli G. (2006) The influence of climate changes in *Platanus* spp. pollination in Spain and Italy. *Grana*, 45: 3, 222-229.
- [63] Hegland S. G., Nielsen A., Lázaro A., Bjerknes A.-S., Totland Ø. (2009) How does climate warming affect plant-pollinator interactions? *Ecology Letters*, 12: 184-195.
- [64] Forrest J. R. K. (2015) Plant - pollinator interactions and phenological change: what can we learn about climate impacts from experiments and observations? *Oikos* 124: 4-13.
- [65] Hua F., Hu J., Liu Y., et. al. (2016) Community-wide changes in intertaxonomic temporal co-occurrence resulting from phenological shifts. *Global Change Biology*, 22, 1746-1754.
- [66] Coristine L. E., Soroye P., Soares R. N., Robillard C., Kerr T. J. (2016) Dispersal Limitation, Climate Change, and Practical Tools for Butterfly Conservation in Intensively Used Landscapes. *Natural Areas Journal*, 36 (4): 440-452.
- [67] Molnár V. A., Tökölly J., Végvári Z., Sramkó G., Sulyok J., Barta Z. (2012) Pollination mode predicts phenological response to climate change in terrestrial orchids: a case study from central Europe. *Journal of Ecology*, 100, 1141-1152.
- [68] Bellard C., Bertelsmeier C., Leadley P., Thuiller W., Courchamp F. (2012) Impacts of climate change on the future of biodiversity. *Ecol Lett.*, 15 (4): 365-377.
- [69] Giannini T. C., Acosta A. L., da Silva C. I., et. al. (2013) Identifying the areas to preserve passion fruit pollination service in Brazilian Tropical Savannas under climate change. *Agriculture, Ecosystems and Environment* 171, 39-46.
- [70] Polce C., Garratt M. P., Termansen M., et. al. (2014) Climate-driven spatial mismatches between British orchards and their pollinators: increased risks of pollination deficits. *Global Change Biology*, 20, 2815-2828.
- [71] Gallai N., Salles J.-M., Settele J., Vaissière B. E. (2009) Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics* 68, 810-821.
- [72] Kaloveloni A., Tscheulin T., Vujic A., Radenkovic S., Petanidou T. (2015) Winners and losers of climate change for the genus *Merodon* (Diptera: Syrphidae) across the Balkan Peninsula. *Ecological Modelling* 313, 201-211.
- [73] Barnosky A. D., Hadly E. A., Gonzalez P., et al. (2017) Merging paleobiology with conservation biology to guide the future of terrestrial ecosystems. *Science* Vol 355, Issue 6325.
- [74] Stein B. A., Glick P., Edelson N., Staudt A. (2014) Climate-Smart Conservation: Putting Adaptation Principles into Practice. National Wildlife Federation, Washington, D. C., (eds.).
- [75] Groves C. R., Game E. T., Anderson M. G., et. al. (2012) Incorporating climate change into systematic conservation planning. *Biodivers Conserv*, 21: 1651-1671.
- [76] Jump A. S., Peñuelas J. (2005) Running to stand still: adaptation and the response of plants to rapid climate change. *Ecology Letters*, 8: 1010-1020.
- [77] Heller N. E., Zavaleta E. S. (2009) Biodiversity management in the face of climate change: A review of 22 years of recommendations. *Biological Conservation* 142, 14-32.
- [78] Davis M. B., Shaw R. G. (2001) Range shifts and adaptive responses to quaternary climate change. *Science*, 292, 5517; ProQuest pg. 673.
- [79] Hunter M. L. Jr., Hutchinson A. (1994) The Virtues and Shortcomings of Parochialism: Conserving Species That Are Locally Rare, but Globally Common. *Conservation Biology*, Vol. 8, No. 4, pp. 1163-1165.
- [80] Olson D. M., Dinerstein E. (1998) The Global 200: A Representation Approach to Conserving the Earth's Most Biologically Valuable Ecoregions. *Conservation Biology*, Vol. 12, No. 3, pp. 502-515.
- [81] Brooks T. M., Mittermeier R. A., da Fonseca G. A. B., et. al. (2006) Global Biodiversity Conservation Priorities. *Science*, New Series, Vol. 313, No. 5783, pp. 58-61.
- [82] Millennium Ecosystem Assessment (2005) Ecosystems and human well-being: synthesis. Island Press, Washington, D. C.
- [83] Lashof D. A., DeAngelo B. J. (1997) Terrestrial Ecosystem Feedbacks to Global Climate Change. *Annu. Rev. Energy Environ.* 22: 75-118.
- [84] Joppa L. N., Loarie S. R., Pimm S. L. (2008) On the protection of "protected areas". *Proc. Natl. Acad. Sci. U.S.A.* 105, 6673-6678.
- [85] Cronk Q. C. B. (1997) Islands: stability, diversity, conservation. *Biodiversity and Conservation* 6, 477-493.

- [86] Wright S. D., Gillman L. N., Ross H. A., Keeling D. J. (2009) Slower Tempo of Microevolution in Island Birds: Implications for Conservation Biology. *Evolution*, 63 (9): 2275-2287.
- [87] Pereira H. M., Navarro L. M., Martins I. S. (2012) Global Biodiversity Change: The Bad, the Good, and the Unknown. *The Annual Review of Environment and Resources*, 37: 25-50.
- [88] Soepadmo E. (1993) TROPICAL RAIN FORESTS AS CARBON SINKS. *Chemosphere*, Vol 27, No. 6, pp 1025-1039.
- [89] Reijnders L., Huijbregts M. A. J. (2008) Palm oil and the emission of carbon-based greenhouse gases. *Journal of Cleaner Production* 16, 477-482.
- [90] Pereira H. M., Leadley P. W., Proença V., et al. (2010) Scenarios for Global Biodiversity in the 21st Century. *Science* Vol 330.
- [91] Tilman D., Fargione J., Wolff B., et. al. (2001) Forecasting Agriculturally Driven Global Environmental Change. *Science* Vol 292.
- [92] Myers N., Kent J. (2003) New Consumers: The Influence of Affluence on the Environment. *Proceedings of the National Academy of Sciences of the United States of America*, Vol 100, No. 8, pp. 4963-4968.
- [93] Pimentel D., Pimentel M. (2003) Sustainability of meat-based and plant-based diets and the environment. *Am J Clin Nutr*, 78 (suppl): 660S-3S.
- [94] Matthews E., Hammond A. (1999) Critical Consumption Trends and Implications: Degrading Earth's Ecosystems. World Resources Institute, Washington, DC.
- [95] Nandwani S. S. (1996) Solar cookers cheap technology with high ecological benefits. *Ecological Economics* 17, 73-81.
- [96] Dravid M. N., Chandak A., Phute S. U., Khadse R. K., Adchitre H. R., Kulkarni S. D. (2012) The use of solar energy for powering a portable autoclave. *Journal of Hospital Infection* 80, 345-347.
- [97] United Nations in India (2015) <http://in.one.un.org/task-teams/sustainable-energy-for-all/>, 14/5-2017, 4.35 p.m.
- [98] Adeney J. M., Christensen N. L., Pimm S. L. (2009) Reserves protect against deforestation fires in the Amazon. *PLOS ONE* 4, e5014.
- [99] Maxwell S. L., Rhodes J. R., Runge M. C., et. al. (2015) How much is new information worth? Evaluating the financial benefit of resolving management uncertainty. *Journal of Applied Ecology*, 52, 12-20.
- [100] Butt N., Possingham H. P., De Los Rios C., et. al. (2016a) Challenges in assessing the vulnerability of species to climate change to inform conservation actions. *Biological Conservation* 199, 10-15.
- [101] Griffith B., Scott J. M., Carpenter J. W., Redd C. (1989) Translocation as a Species Conservation Tool: Status and Strategy. *Science*; 245, 4917; ProQuest pg. 477.
- [102] Millar C. I., Stephenson N. L., Stephens S. L. (2007) Climate Change and Forests of the Future: Managing in the Face of Uncertainty. *Ecological Applications*, Vol. 17, No. 8, pp. 2145-2151.
- [103] Barnosky A. D. (2009) Managing nature as Earth warms. *Nature* Vol 458.
- [104] Tricarico E. (2016) Do alien invasive species and climate change foster conservation behaviour? *Aquatic Conserv: Mar. Freshw. Ecosyst.* 26: 228-232.
- [105] Butt N., Whiting S., Dethmers K. (2016b) Identifying future sea turtle conservation areas under climate change. *Biological Conservation* 204, 189-196.
- [106] Kleiman D. G. (1989) Reintroduction of Captive Mammals for Conservation. *BioScience*, Vol. 39, No. 3, pp. 152-161.
- [107] Hunter M. L. Jr (2007) Climate Change and Moving Species: Furthering the Debate on Assisted Colonization. *Conservation Biology* Volume 21, No. 5, 1356-1358.
- [108] Payne B. L., Bro-Jørgensen J. (2016) A framework for prioritizing conservation translocations to mimic natural ecological processes under climate change: A case study with African antelopes. *Biological Conservation* 201, 230-236.