



Efficacy of Herbicides to Various Weeds in Perennial Crops

Bogale Ayana

Ethiopian Institute of Agricultural Research, Holeta Agricultural Research Center, Holeta, Ethiopia

Email address:

bogalayana@gmail.com

To cite this article:

Bogale Ayana. Efficacy of Herbicides to Various Weeds in Perennial Crops. *American Journal of Chemical and Biochemical Engineering*. Vol. 6, No. 1, 2022, pp. 1-5. doi: 10.11648/j.ajcbe.20220601.11

Received: February 25, 2022; **Accepted:** April 12, 2022; **Published:** April 20, 2022

Abstract: Peach is one of the most important fruit crops in the world with wider ecological adaptation which has been cultivated in sub-tropical to temperate climates. In spite of its economic importance, its production has been affected by various abiotic and biotic factors. Therefore, the objective of this study was to assess the effect of non-selective herbicides against peach weeds. Treatments consisted of three post emergence non selective herbicides; Glynosh @ 1.75kg/ha, Glynosh @ 1.75 L ha⁻¹, Glyphosate @ 2L ha⁻¹ along with weedy check that were laid out in RCBD with three replications. The field was infested with nine weed species in which five species were annual weeds, three species were perennial weeds and one species was under category of biennial. The maximum relative weed density (17.38) was calculated from *Cynodon dactylon* L. while minimum (3.90) number was observed from *Centella asiatica* L. The result also showed that application of all herbicides had no statistically significant differences on *Bidens pilosa* L., *Medicago polymorpha* L., *Palntago lanceolata* L. and *Raphanus raphanistrum* L. except for weedy check. In addition, application of Glyphosate IPA 480gm./lit, 48% SL and Glyphosate produced statistically no significant differences on the remaining weed species. Application of all herbicides revealed statistically no significant differences in terms of weed dry weight while better weed control efficiency and fruit yield was obtained from application of Glyphosate IPA 480gm./lit, 48% SL at all experimental sites. Furthermore, there is no phytotoxicity was observed due to candidate herbicide if applied with great care in between row planted peach. Thus, Glynosh 1.75 L/ha is recommended for control of various annual and perennial weeds in peach.

Keywords: Difference, Effect, Glynosh, Herbicides, Weed

1. Introduction

Peach is one of the most important fruit crops in the world, with wider ecological adaptation, cultivated in sub-tropical to temperate climates. It is cultivated on 1.54 million ha with an annual production of 20.27 million tons in the world [1]. The broad wider use of peach including its importance in a healthy and balanced diet has resulted in increased production of both processed and fresh varieties worldwide in the recent years. Similarly, a marked increase in area of production has also been observed in recent years in Ethiopia. The crop is among grown fruit crops using rain fed both by smallholder and commercial farmers in highlands of Ethiopia.

According to reports of [2, 3] revealed that weed interference has been reported to affect tree growth, yield, and fruit quality in peach. However, the magnitude of the effect on fruit yield and size depends on the weed species [4]. Cultivation or herbicides are most commonly used for the management of weeds in peach

orchards. Weed control by cultivation with discs and cultivators is temporary and not cost effective. Further, cultivation with discs and cultivators cannot manage the weeds under the trees in between the tree rows; hence, manual weeding is performed or herbicides are used. High labor costs in developed nations make manual weed in uneconomical. Conventional agricultural practices rely on synthetic herbicides for managing weeds, and these compounds account for more than half of the volume of all agricultural pesticides applied in the developed world [5]. The increased pressure to reduce herbicide applications and new interests in organic farming underline the importance of alternative approaches for orchard weed suppression [6]. Further, repeated use of the same combination of herbicides can result in a shift in weed population and development of resistance. Rotation of herbicides with a different mode of action avoids the problem of weed resistance and improves weed control.

In spite of the crop's wider ecological adaptation, both biotic and abiotic stresses had limitation on its production and productivity. Weeds control is less noticeable to Peach

production in Ethiopia and economic losses resulting from crop damage or weed control measures are less noticeable.

In Ethiopia, weeds are more severe during summer when rainfall is high. Weeds infested quickly in fields and can result in total crop failure if left untreated. Sustainable control of weeds actually requires integration of cultural practices that involves the application of herbicides. However, due to different germination time of the weeds, herbicides with better efficacy should be made available for the growers to reduce the consequential yield loss caused by the weeds. Therefore, the objective of this study was to assess the effect of non-selective herbicides against peach weeds.

2. Materials and Methods

2.1. Description of the Study Area

Field experiment was conducted from June, 2020 to December, 2020 main cropping season under rain fed conditions at Holeta Agricultural Research Center and Medegudina locations. Holeta is located 33km west of Addis Ababa at an elevation of 2400 m.a.s.l and within the geographic coordinates of 9° 00'N and 38° 30'E. The area receives annual rain fall of 1144 mm with mean minimum and maximum temperatures of 6°C and 22°C respectively (EIAR, b2018). The soil of the experimental field is clay loam with pH of 6.65, organic carbon (2.26%), available Phosphorus (14.17 mgkg⁻¹), total nitrogen (0.12%) and cation exchange capacity of (17Cmol/kg⁻¹) (EIAR, 2018). The edaphic and climatic conditions observed during the trial period were favorable for the exuberant growth of numerous weed species that competed with the crop plants. The climatic conditions observed during the trial period mean rain fall of 1114.5 mm relative humidity 78.8% with mean minimum and maximum temperatures of 8°C and 25.2°C, respectively. The remaining area received almost similar weather conditions with that of Holeta.

2.2. Treatments and Crop Management

Field experiment was conducted at Holeta Agricultural Research Center and Medegudina farmer field where fields were infested with many weed species. Treatments were laid out plot sizes of 4m×4m along with the test herbicides; Glyphosh 1.75 kg/ha, Glyphosh 1.75 L/ha, Glyphosate 2 L/ha

and weedy check. The design was RCBD with three replications. Peaches were already planted at the experimental sites at recommended spacing. All agronomic practices were applied based on recommendation for peach. All herbicides were applied as post emergence with spray volume of 250L/ha.

2.3. Data Collection

Weed species identification was made by uprooting fresh weeds from experimental field and taken to laboratory. After the weed flora were identified, they were categorized as grasses and broad leaf weeds using reference of manuals, consulting experienced professionals and comparing with existed herbarium as described by [7]. Data regarding the kind of weed species and their densities were counted at 25 day after sowing i. e, before the application of herbicides by using four quadrats with sizes of 0.25 m x 0.25 m randomly placed in each plot and their density was calculated m⁻². In addition, individual weed species density count was also done at 45 days after herbicides were applied by randomly placing four quadrats of sizes 0.25m×0.25m converted to m⁻². The density of each weed species in the field was counted after treatment by randomly placing of four quadrat of sizes 25 cm×25 cm in each plots and calculated m⁻² basis. The relative weed density was calculated by the formula [8]: $RWD (\%) = \frac{\text{number of weeds of species}}{\text{total weeds of species}} \times 100$

The aboveground dry biomass of grass weeds and dry biomass of broad leaf weeds harvested from each quadrat placed into paper bags separately and oven drying at a 65°C for 48 hours and subsequently the dry weights were measured. Weed control efficiency (WCE) was determined by the following formula $WCE (\%) = \frac{WDC - DWP}{WDC} \times 100$, where, WCE=Weed Control Efficiency, WDC=Weed Dry weight in Control Plot and DWP = Weed Dry weight in Particular treatment [9]. Grain yield was measured after threshing the sun dried plants harvested from each net plot and the yield was adjusted at 12.5% grain moisture content [10].

2.4. Statistical Analysis

The means of each data was checked by the normality test depending on Shapiro test (Pr < W) before analysis of variance using the GLM procedure of SAS (SAS 9.3 version). When the treatment effects were significant, means were compared using Fisher's LSD test at 5% level of significance [11].

Table 1. Trade name, common name, rates and mode of action of herbicides.

Trade name	Common name	Rate/ha	Selectivity
Glynosh	Glyphosate IPA 75.7% WSG	1.75 kg/ha	Non selective
Glynosh	Glyphosate IPA 480gm./lit, 48% SL	1.75 L/ha	Non selective
Round up	Glyphosate	2 L/ha	Non selective
Weedy check	-	-	-

3. Results and Discussions

3.1. Identification of Weed Flora in the Experimental Field

The experimental sites were infested with various weed floras that are problematic in perennial crops as well as

pastures. Nine weed species were identified from the experimental location in which five species categorized as annual, three species were under category of perennial and only one species under biennial (Table 2). This result showed that the field was highly infested with annual weeds. The maximum relative weed density (17.38%) was calculated from *C. dactylon* L. while minimum (3.90%) number was

observed from *C. asiatica* L. which indicated that perennial grasses are more problematic weed in perennial crop (Peach).

Table 2. Weed species, relative density and life form in the experimental field.

Scientific names	Families	Weed density count m ² before spray	Relative weed density (%)	Life form
<i>Bidens plosa</i> L.	Asteraceae	460	15.70	Annual
<i>Ocimum australis</i> L.	Lamiaceae	336	11.47	Biennial
<i>Medicago polymorpha</i> L.	Fabaceae	420	14.34	Perennial
<i>Cynodon dactylon</i> L.	Poaceae	509	17.38	Perennial
<i>Bidens pachyloma</i> L.	Asteraceae	372	12.70	Annual
<i>Plantago lanceolata</i> L.	Plantaginaceae	132	4.50	Annual
<i>Centella asiatica</i> L.	Apiaceae	116	3.90	Annual
<i>Rumex abyssinicus</i> Jacq	Polygonaceae	204	6.97	Perennial
<i>Raphanus raphanistrum</i> L.	Brassicaceae	380	12.97	Annual

3.2. Weed Density Count at 45 Days After Herbicides Application at Holeta

Individual weed densities were significantly affected by all tested herbicides in peach (Table 3). The result showed that application of all herbicides had no statistically significant differences on *B. pilosa*, *M. polymorpha*, *P. lanceolata* and *R. raphanistrum* except for weedy check. Similarly, application

of Glyphosate IPA 480gm./lit, 48% SL and Glyphosate produced statistically no significant differences on *Ocimum spp.*, *M. polymorpha*, *B. pachyloma*, *P. lanceolata*, *C. asiatica* and *R. raphanistrum*. This result revealed that herbicides having the same mode of action had similar performance on targeted weed species. On other hand, broad spectrum herbicides have the ability of controlling of various weed species that could be more preferable for farmers.

Table 3. Effects of Herbicides on individual weed species at 45 days after herbicides application at Holeta.

Treatments	<i>Bidens pilosa</i> L.	<i>Ocimum spp.</i> L.	<i>Medicago polymorpha</i> L.	<i>Cynodon dactylon</i> L.	<i>Bidens pachyloma</i> L.	<i>Palntago lanceolata</i> L.	<i>Centella asiatica</i> L.	<i>Rumex abyssinica</i> Jacq	<i>Raphanus raphanistrum</i> L.
Glyphosate IPA 75.7% WSG	11.33b	12.00b	16.00b	173.00ab	14.00b	10.00b	14.66b	13.00b	14.00b
Glyphosate IPA 480gm./lit, 48% SL	3.33b	4.00c	4.00b	148.00ab	4.00c	4.00b	4.00c	4.00c	4.00b
Glyphosate	3.33b	4.00c	4.00b	119.00b	4.00c	4.00b	4.00c	4.00c	4.00b
Weedy check	30.66a	14.66a	45.87a	184.00a	20.67a	16.00a	77.33a	28.00a	18.66a
LSD (5%)	15.35	2.31	24.87	64.52	3.12	4.76	8.29	7.88	7.56
CV (%)	6.3	13	21	20	14.57	23.80	16.59	32.22	33.90

3.3. Weed Density Count at 45 Days After Herbicides Application at Medegudina

Individual weed densities were significantly affected by all tested herbicides in peach (Table 4). The result showed that application of all herbicides had no statistically significant differences on *B. pilosa*, *M. polymorpha*, *P. lanceolata* and *R. raphanistrum* except for weedy check. Similarly, application

of Glyphosate IPA 480gm./lit, 48% SL and Glyphosate produced statistically no significant differences on *Ocimum spp.*, *M. polymorpha*, *B. pachyloma*, *P. lanceolata*, *C. asiatica* and *R. raphanistrum*. This result revealed that herbicides having the same mode of action had similar performance on targeted weed species. On other hand, broad spectrum herbicides have the ability of controlling of various weed species that could be more preferable for farmers.

Table 4. Effects of Herbicides on individual weed species at 45 days after herbicides application at Medegudina.

Treatments	<i>Bidens pilosa</i> spp.	<i>Ocimum spp.</i>	<i>Medicago polymorpha</i>	<i>Cynodon dactylon</i>	<i>Bidens pachyloma</i>	<i>Palntago lanceolata</i>	<i>Centella asiatica</i>	<i>Rumex abyssinica</i>	<i>Raphanus raphanistrum</i>
Glyphosate IPA 75.7% WSG	13.33b	14.00b	18.00b	175.00ab	14.00b	10.00b	14.66b	13.00b	14.00b
Glyphosate IPA 480gm./lit, 48% SL	5.33b	6.00c	6.00b	150.33ab	4.00c	4.00b	4.00c	4.00c	4.00b
Glyphosate	5.33b	6.00c	6.00b	121.33b	4.00c	4.00b	4.00c	4.00c	4.00b
Weedy check	32.66a	16.66a	47.33a	186.00a	20.67a	16.00a	77.33a	28.00a	18.66a
LSD (5%)	15.34	2.30	24.87	64.52	3.10	4.75	8.29	7.88	7.56
CV (%)	5.4	10.82	6.4	20.41	12.27	19.83	15.37	27.70	28.75

In general, results of both locations produced almost similar results of herbicidal treatment in terms of weed control. Herbicides having the same mode of action may produce similar results that could be better advantages for

peach growers.

3.4. Weed Dry Weight

Weed dry weight was significantly affected by

application of different nonselective herbicides (Table 5). There was no dry weed biomass was recorded from application of Weed free plots at both locations while the maximum weed dry weight (2633.33 kg/ha) at Holeta and (2433.33kg/ha) at Medegudina were recorded from weedy check plots. There is no statistically significant differences were observed due to application of all herbicides at both locations except for weedy check. Barley that the lowest dry weight recorded was due to removal of most of the weed plants there which suppressed density of weeds and resulting into a lower competition between the crop and weeds for resources [12]. This finding was also similarly with [13, 14] who concluded that maximum weed density in weedy check can be attributed to unchecked growth, while application of herbicide caused mortality of weed resulting in lower weed density at harvest.

3.5. Weed Control Efficiency

Weed control efficiency was significantly affected by application of different herbicides (Table 5). The maximum weed control efficiency (95.88%) at Holeta and (96.75%) at Medegudina was recorded from application of Glyphosate IPA 480gm./lit, 48% SL while no weed control efficiency at weedy check plots at all tested locations. The maximum weed control efficiency from application of Weed free plots

due to minimum weed dry biomass. This result was similar with [12] who stated that the reduction in weed dry weight in barley might be due to the inhibition effect of treatments on growth and development of weeds. This finding is also similar with [15, 16] who reported that the high control efficiency indicated that the weed were controlled when they are young or before they accumulated more dry matter by competing with the crop plants.

3.6. Marketable Fruit Yield

Fruit yield was significantly affected by application of different herbicides (Table 5). The maximum yield (26.3tons/ha) was recorded from application of Glyphosate IPA 480gm./lit, 48% SL at Holeta while the minimum values (1.0 ton/ha) was recorded from weedy check at Holeta. In addition, there is no statistically significant variation was observed due to application of all herbicides except for weedy check at Medegudina. The maximum fruit yield from application of Glyphosate IPA 480gm./lit, 48% SL implied that better weed control that enable the plants to utilize more growth resources but the minimum fruit yield at weedy check probably due to severe competitions of weeds. This result was analogous with [17] who stated that the maximum grain yield was obtained where minimum weed crop competition for nutrients and water was existed.

Table 5. Effects of Herbicides application weed dry weight, weed control efficiency and yield at Holeta and Medegudina.

Herbicides	Weed dry weight (kg/ha)		Weed control efficiency (%)		Yield (ton/ha)	
	Holeta	Medegudina	Holeta	Medegudina	Holeta	Medegudina
Glyphosate IPA 75.7% WSG	153.33b	123.33b	94.17b	95.54b	23.0b	21.3b
Glyphosate IPA 480gm./lit, 48% SL	108.33b	90.00b	95.88a	96.75a	26.3a	25.0b
Glyphosate	200.00b	170.00b	92.40c	93.85c	23.6b	22.0b
Weedy check	2633.33a	2433.33a	0.00d	0.00d	1.0c	0.80c
LSD (5%)	135.72	135.30	0.90	1.07	2.20	2.23
CV (%)	8.77	9.61	0.64	0.75	5.32	5.85

4. Conclusion

Peach is one of the most important fruit crops in the world with wider ecological adaptation which has been cultivated in sub-tropical to temperate climates. In spite of its economic importance, its production has been affected by various abiotic and biotic factors. Among biotic factors weeds can share greatest yield reducing factors if left uncontrolled. Hence, the aim of this study was to test the efficacy non selective herbicides against peach weeds. The experimental sites were infested with various weed floras that are problematic in perennial crops as well as pastures. The maximum relative weed density was calculated from *C. dactylon* L. while minimum number was observed from *C. asiatica*. Individual weed densities were significantly affected by all tested herbicides in peach. The result also showed that application of all herbicides had no statistically significant differences on *B. pilosa*, *M. polymorpha*, *P. lanceolata* and *R. raphanistrum* except for weedy check. Correspondingly, application of Glyphosate IPA 480gm./lit,

48% SL and Glyphosate produced statistically no significant differences on *Ocimum spp.*, *M. polymorpha*, *B. pachyloma*, *P. lanceolata*, *C. asiatica* and *R. raphanistrum*. The result revealed that maximum weed control efficacy was recorded from application of Glynosh (Glyphosate IPA 480gm./lit, 48% SL) than other tested herbicides. Application of all herbicides revealed statistically no significant differences in terms of weed dry weight while better weed control efficiency and fruit yield was obtained from application of Glyphosate IPA 480gm./lit, 48% SL at all experimental sites. Furthermore, there is no phytotoxicity was observed due to herbicides if applied with great care in between row planted peach. Thus, Glynosh (Glyphosate IPA 480gm./lit, 48% SL) at the rate of 1.75L/ha in 250 L of water per hectare is recommended as alternative herbicide for control of various annual and perennial weeds in peach.

Declaration of Competing Interest

The author declared that there were no competing interests.

References

- [1] FAOSTAT, 2010. Statistical database of the food and agricultural organization of the United Nations [Internet]. [cited 2012 December 15]. Available from: <http://www.faostat.fao.org>
- [2] Majek, B. A., Neary, P. E., Polk, D. F., 1993. Smooth pigweed interference in newly planted peach trees. *Journal of Product Agriculture*. 6, 244-246.
- [3] Parker, M., Meyer, J. R., 1996. Peach tree vegetative and root growth respond to orchard floor management. *Horticultural Science*. 3 (1), 330-333.
- [4] Tworkoski, T. J., Glenn, D. M., 2001. Yield, shoots and root growth, and physiological responses of mature peach trees to grass competition. *Horticultural Science*. 3 (6), 1214-1218.
- [5] Dayan, F. E, Howell, J., Marais J. P, Ferreira, D., Koivunen, M., 2011. Manuka oil, a natural herbicide with pre emergence activity. *Weed Science*. 59, 464-469.
- [6] Goh, K. M., Pearson, D. R., Daly, M. J., 2001. Effects of apple orchard production systems on some important soil physical, chemical and biological quality parameters. *Biol. Agric. Hortic*. 18, 269-292.
- [7] Stroud, A. and Parker, C., 1989. A weed identification guide for Ethiopia. *A weed identification guide for Ethiopia*: pp. 55-78.
- [8] Marwat, K. S., Usman, K., Khan, N., Khan, U. M., Ahmad, E., Khan, A. E., Khan, A. M., Rehman, U. A., 2013. Weeds of Wheat Crop and Their Control Strategies in Dera Ismail Khan District, Khyber Pakhtun Khwa, Pakistan. *American Journal of Plant Sciences*. 4, 66-76.
- [9] Davasenapathy PT, Remesh B., 2008. Efficiency indices for Agricultural Management Research. New Indian Publishing Agency, New Delhi India: 576-64.
- [10] Amare, T., Sharma, J. J. and Kassahun, Z., 2014. "Effect of Weed Control Methods on Weeds and Wheat (*Triticum Aestivum* L.) Yield." *World Journal of Agricultural Research*. vol. 2, no. 3 (2014): 124-128.
- [11] Gomez, K. A. and Gomez, A. A., 1984. *Statistical procedures for agricultural research* (2 ed.). John Wiley and sons, New York, pp. 680.
- [12] Megersa K, Tigist B, Geleta G, Girma C, Kasa M, Megersa D, Hailu F., 2017. Effect of Various Weed Management Options on Weeds and Yield of Barley (*Hordeum vulgare* L.) at Shambo and Gedo, Western Oromia. *Journal of Biology, Agriculture and Healthcare*, 7 (21): 74-83.
- [13] Munsif, F., Ali, K., Khan, I., Khan, H. U. and Anwar, M., 2009. Efficacy of various herbicides against weeds and their impact on yield of maize. *Pak. J. Weed Sci. Res*, 15 (2-3), pp. 191-198.
- [14] Nadeem, M. A., Ali, A. and Tanveer, A. S. I. F., 2006. Effect of different weed control practices and fertilizer levels on the weeds and grain yield of wheat. *Pakistan Journal of Botany*, 39 (1), p. 173.
- [15] Ashiq, Muhammad, N. and Ahmad, N., 2007. Comparative efficacy of different herbicides against broadleaved weeds in wheat. *Pakistan Journal of Weed Science Research*, 13 (3-4): pp. 149-156.
- [16] Tana, T., Lemlem, M. and Dejene, M., 2018. Determination of Critical Period of Weed Competition in Bread Wheat *Triticum aestivum* L. at Haramaya, Eastern Ethiopia, Doctoral dissertation, Haramaya University. pp. 1-92.
- [17] Shah A M, Ali S, Ahmad I, Wazir G, Shafique O, Hanif MA, Khan BA and Zareen S., 2018. Weeds population studies and wheat productivity as influenced by different sowing techniques and herbicides. *Pakistan Journal of Agricultural Research*, 32 (1): 87-94.