

Nutrient Exports Under Different Harvesting Regimes in Two Types of Larch Plantation with Different Age in Northeastern China

Hongfeng Wang^{1,2}, Xueyun Dong³, Yajuan Xing^{2,4}, Zhengquan Wang¹, Guoyong Yan^{1,2}, Jianyu Wang^{1,2}, Qinggui Wang^{1,2,*}

¹School of Forestry, Northeast Forestry University, Harbin, China

²College of Agricultural Resource and Environment, Heilongjiang University, Harbin, China

³School of Science, Harbin University, Harbin, China

⁴Institute of Forestry Science of Heilongjiang Province, Harbin, China

Email address:

qgwang1970@163.com (Qinggui Wang)

*Corresponding author

To cite this article:

Hongfeng Wang, Xueyun Dong, Yajuan Xing, Zhengquan Wang, Guoyong Yan, Jianyu Wang, Qinggui Wang. Nutrient Exports Under Different Harvesting Regimes in Two Types of Larch Plantation with Different Age in Northeastern China. *Journal of Energy and Natural Resources*. Vol. 5, No. 6, 2016, pp. 67-77. doi: 10.11648/j.jenr.20160506.11

Received: October 29, 2016; **Accepted:** November 29, 2016; **Published:** December 29, 2016

Abstract: It has been confirmed in studies conducted in different regions and forest types that the harvesting regime has a significant impact on the export of nutrients from a plantation. Through establishing allometric growth equations and studying nutrient distribution patterns, the amounts of nutrients exported from larch (*Larix* spp.) plantations with different stand ages under different harvesting regimes (removal of stem wood, branches and bark (Rsbb), removal of stem wood and bark (Rsb), and removal of stem wood (Rs)) were determined in the present study. The results showed the following: 1) the organs of 20-year-old and 40-year-old larch trees exhibited basically the same nutrient distribution pattern. However, due to the difference in biomass partitioning in trees of these ages, they showed a significant difference in the cumulative amounts of nutrients in their organs. Specifically, the cumulative amounts of nutrient elements in the leaves and old branches of 20-year-old larch trees were significantly higher than in 40-year-old larch trees. However, the cumulative amounts of various nutrients in the xylem of the trunks of 20-year-old larch trees were significantly lower than in 40-year-old larch trees. 2) Compared with the Rsbb20, Rsbb40 or Rsb20 could result in significant decreases in the annual mean amounts of nutrients exported, the ratio of the annual mean total amount of nutrients exported to the annual net cumulative total amount of nutrients and the proportion of soil nutrients in the 0-30 cm soil layer. Changing the Rsbb20 to an Rsb40 or an Rs40 could further reduce the annual mean amounts of nutrients exported, the ratio of the annual mean total amount of nutrients exported to the annual net cumulative total amount of nutrients and the proportion of soil nutrients in the 0-30 cm soil layer. 3) Among the examined harvesting regimes, the Rs20 resulted in the lowest annual mean amounts of nutrients being exported. However, there was no significant difference in the annual mean amounts of nutrients exported between the Rs20 and the Rs40. The amounts of N, Ca and Mg exported under the Rs20 were slightly higher than under the Rs40, whereas the amounts of P and K exported under the Rs20 were slightly lower than under the Rs40. Hence, the harvesting regime is an important factor that results in the export of system nutrients and a decline in soil fertility. Therefore, prolonging the harvesting cycle and adopting Rs are two options for reducing the nutrients export.

Keywords: Allometric Growth Equation, Different Stand Ages, Larch Plantation, Biomass, Nutrient Distribution Pattern

1. Introduction

Much attention has long been paid to the decline in soil

fertility in plantations [1], [2]. Maintaining a balance between the import and export of nutrient elements to ensure the stability of the nutrient capital of the soil is the basis for a

forest ecosystem to achieve sustainable productivity [3]. We often observe this type of balance in natural ecosystems [4]. However, the periodic harvesting of timber may result in disruption of the balance between the import and export of nutrient elements in plantations, which account for 7% of the total forest area worldwide [5]. Researchers have indeed observed a decline in soil nutrients caused by harvesting in different types of plantations [6], [7], [8], [9]. Therefore, how to maintain the capacity of a soil to provide a long-term nutrient supply under continued harvesting has been attracting increasing attention [10].

Through numerous studies involving allometric growth equations and nutrient distribution patterns, it has been found that different harvesting regimes have a significant impact on the maintenance of the capacity of a soil to provide a long-term nutrient supply [6], [9]. On this basis, government agencies have formulated regulations and various rules to ensure the sustainable utilization of nutrients in plantations. For example, current guidelines by Wisconsin Department of Natural Resources (WDNRs) for the harvesting of forested areas recommend that detritus with a diameter less than 10cm be left on-site along with roots, stumps, the litter layer, and any woody detritus that was already on the forest floor [11]. States from Maine to Missouri, Canada, and some European countries also made the similar rules [12].

While the importance of harvesting in the cycling of nutrients in plantation ecosystems has been acknowledged [10], [13] and researchers have carried out many studies on the nutrient utilization rates of different tree species [14], [15], [16], [17], [18], there are significant differences between different regions and different tree species [19]. One study showed that a decline in soil fertility can easily occur in fast-growing, high-yield plantations in tropical regions [20], whereas declines in soil fertility do not readily occur in temperate forests with abundant nutrients, such as the forest areas in northeastern China [21]. However, a recent study revealed an increasingly prominent declining trend in soil fertility in the larch plantations in northeastern China [22], [23], [24], [25], [26], [27]. The forest areas in northeastern China account for 31.4% of the total forest area in China. The area of natural forest in the forest areas of northeastern China is gradually decreasing, whereas the plantation area in these forest areas is continuously increasing. Larch (*Larix spp.*) is one of the main timber species in the forest areas in northeastern China. Large areas of pure larch forests have been created since the 1950s. The larch plantation area currently covers 2.0×10^6 hm², accounting for approximately 55% of the total plantation area in northeastern China [28]. If the declining trend in soil fertility continues, a serious threat will be posed to the plantations in northeastern China, or even the entire country. Studies in other regions and forest types have shown that different harvesting regimes (e.g., different harvesting cycles or harvesting regimes involving the removal of stem wood, branches and bark (Rsbb) or only stem wood and bark (Rsb)) have a significant impact on the export of nutrients from plantations [6], [29], [30], [31]. However, we still do not understand the impact of different harvesting

regimes on the export of nutrients from the larch plantations in northeastern China and whether the harvesting regime has already become the chief cause of declining soil fertility in this type of plantation.

Studying nutrient distribution patterns can provide accurate information for estimation of the loss of nutrients caused by harvesting [6], [9]. Through establishing allometric growth equations and investigating nutrient distribution patterns, the impact of different harvesting regimes (Rsbb, Rsb, removal of stem wood (Rs)) on the export of nutrients from larch plantations with two different stand ages was evaluated in the present study, and whether the difference in the export of nutrients would impact the soil nutrient contents and result in a decline in soil fertility in the plantations was assessed. Finally, the impacts of different harvesting regimes on plantation site conditions were determined, providing a basis for scientifically selecting a harvesting regime in northeastern China.

2. Materials and Methods

2.1. Site Characteristics

The study was conducted at the Maoershan Experimental Station (45°21'–45°25'N, 127°30'–127°34'E) of the Northeast Forestry University in northeastern China. The site has a continental temperate monsoon climate with the mean January, July, and annual air temperatures of −19.6°C, 20.9°C, and 2.8 °C, respectively. The mean annual precipitation is 723mm with 477 mm falling from June to August. The soils are Hap-Boric Luvisols, well drained with high organic matter and frozen to a depth of 1m during the winter (from December to April).

2.2. Biomass Determination

In March 2012, three experimental plantation plots (20 m×20 m) with a stand age of 20 years and three experimental plantation plots (20 m×20 m) with a stand age of 40 years were selected at the Maoershan Experimental Station of the Northeast Forestry University. In each plot, five standard trees were selected and cut down. The biomass of the trunk, branches and leaves of each standard tree was determined using the vertical stratifying method, while the biomass of the roots of each standard tree was determined using the whole digging method [32]. The samples collected from different parts of each standard tree were dried, after which the biomass of each organ was calculated. Then, a correlation model between the biomass of each organ and the diameter at breast height (DBH) was established.

2.3. Nutrient Contents in Different Parts of the Trees and Soil

Samples were collected from the trunks, bark, leaves, branches and roots of standard larch trees in stands with different ages. After the samples were ground and sieved, nutrient contents were determined. C and N contents were determined using an element analyzer after digesting the

samples with concentrated $\text{H}_2\text{SO}_4\text{--H}_2\text{O}_2$ (Elementary Vario EL III, Germany). The P content was determined using the Mo-Sb colorimetric method [33]. The digestion solution was also employed for the determination of K, Ca and Mg contents (Atomic Absorption Spectroscopy, PE Aanalyst 800, USA) using the atomic absorption method [34], [35], [36].

Soil samples collected from the field were dried in air, after which the soil nutrient contents were determined. The total N content was determined using the Kjeldahl method. The total P content was determined using the $\text{HClO}_4\text{--H}_2\text{SO}_4$ method. The total K content was determined using the HF--HClO_4 digestion method, and Ca^{2+} and Mg^{2+} contents were determined using the atomic absorption method. The cumulative amount of each soil nutrient was the product of the soil nutrient content and the soil volume [37].

2.4. Statistical Analyses

The cumulative amount of each nutrient in each organ of a single tree was the product of the concentration of the nutrient and the biomass of the organ [38]. The cumulative total amount of nutrients in each organ was the sum of the cumulative amounts of nutrients in the sections at different heights. The biomass, cumulative amounts of nutrients and C storage at the stand level were obtained by multiplying the relevant data for a standard tree by the stand density. Statistical data analysis was performed with SPSS 13.0. The prediction results of the biomass optimization model for each organ were examined using the mean accuracy (average accuracy, AA):

$$AA = 1 - \sum (|a - e| / a) / N \times 100\% \quad (1)$$

where a represents the true value, and e represents the estimated value. The closer the value of AA is to 1, the better the prediction results of the model are. Generally, AA was $>80\%$, indicating that the model was relatively consistent with the actual conditions [39], [40]. The significance of the difference was compared using the least significant difference method ($\alpha=0.05$).

The amounts of nutrients exported under different harvesting regimes were estimated based on the biomass model and the cumulative amounts of nutrient elements in the organs. The examined harvesting regimes were as follows:

Harvesting regime 1: removal of stem wood, branch and bark (Rsbb), which includes two types, one is Rsbb₂₀ that Rsbb harvesting regime with a 20-year harvesting cycle, another is Rsbb₄₀ that Rsbb harvesting regime with a 40-year harvesting cycle.

Harvesting regime 2: removal of stem wood, and bark (Rsb), which includes two types, one is Rsb₂₀ that Rsb harvesting regime with a 20-year harvesting cycle, another is Rsb₄₀ that Rsb harvesting regime with a 40-year harvesting cycle.

Harvesting regime 3: removal of stem wood (Rs), which includes two types, one is Rs₂₀ that Rs harvesting regime with a 20-year harvesting cycle, another is Rs₄₀ that Rs harvesting regime with a 40-year harvesting cycle.

3. Results

3.1. Soil Properties

Similar to previous research findings [21], the soil in the study area contained relatively high nutrient contents (Table 1). The N, P and K contents of the soil all gradually decreased with increasing soil layer depths, but to different degrees. The N and K contents decreased to a relatively large degree, whereas the P content decreased to a relatively small degree with an increasing soil layer depth. The N content in the 0–10 cm soil layer was 0.963% and decreased to 0.412% in the 21–30 cm soil layer (an over 50% decrease). The K content in the 0–10 cm soil layer was 0.35% and decreased to 0.1% in the 21–30 cm soil layer (an over 70% decrease). There were no significant differences in the Ca and Mg contents between different soil layers. In addition, the Ca and Mg contents did not exhibit a clear increasing or decreasing trend. The soil density gradually increased with increasing soil layer depth. The soil density was 0.62 g/cm³ in the 0–10 cm soil layer and increased to 1.5 g/cm³ in the 21–30 cm soil layer (an over 50% increase).

Table 1. Basic physical and chemical properties of the soil in the larch plantation.

Nutrients element	Soil layers	Soil nutrients (%)
N	0-10cm	0.963
	11-20cm	0.692
	21-30cm	0.412
P	0-10cm	0.159
	11-20cm	0.137
	21-30cm	0.111
K	0-10cm	0.350
	11-20cm	0.250
	21-30cm	0.100
Ca	0-10cm	0.760
	11-20cm	0.630
	21-30cm	0.730
Mg	0-10cm	0.007
	11-20cm	0.017
	21-30cm	0.026
Soil density g/cm ³	0-10cm	0.620
	11-20cm	0.930
	21-30cm	1.500

3.2. Biomass Production and Nutrient Concentrations in Tree Components

A regression equation for DBH and the biomass of each organ was established using 10 standard trees:

$$W = aD^b \quad (2)$$

Where W represents the biomass, and D represents DBH (diameter at breast height).

The coefficient of determination (R^2) and the residual sum of squares (RSS) were used as the evaluation indexes to evaluate the fitting results of each model. The results show that among the 22 biomass models for the 11 indexes (total biomass; aboveground biomass; biomass of the xylem of the trunk; biomass of the bark; biomass of the stalks; biomass of the branches; biomass of the new branches; biomass of the old

branches; biomass of the leaves; belowground biomass; and biomass of the root system) of the larch trees with two different stand ages, 11 exhibited an R^2 between 0.814 and 0.991, an RSS between 0.007 and 0.081 and an evaluation accuracy between 81.7% and 95.1% (Table 2), indicating that the models established with the DBH as an independent variable could satisfactorily predict the biomass of each organ of a larch tree. However, the model for the biomass of the new branches of the 20-year-old larch trees showed an R^2 of only 0.659, an RSS of 0.754 and a mean accuracy of only 40.4%

(Table 2), indicating that this model displayed a low prediction accuracy. The biomass of the new branches of the 20-year-old larch trees was reestablished using the regression equation for DBH and tree height:

$$W = a(D^2H)^b \quad (3)$$

Where W represents the biomass, and D represents DBH (diameter at breast height), and H represents height of tree.

Table 2. Single-tree biomass estimation model for the larch plantation and examination of the accuracy of the model.

Organ	Regression equation	Coefficient		R^2	Sum of squared residuals	Average accuracy (%)
	$M = aD^b$	a	b			
Total	40 years	1.785	2.676	0.925	0.025	91.3
	20 years	1.117	2.607	0.947	0.067	84.3
Above ground	40 years	1.555	2.826	0.908	0.034	88.7
	20 years	1.051	2.593	0.945	0.069	84.1
Woody tissues of Stem	40 years	1.704	2.685	0.931	0.023	91.2
	20 years	1.700	2.416	0.982	0.019	92.9
Bark	40 years	0.889	2.528	0.814	0.061	84.7
	20 years	0.886	2.512	0.935	0.077	81.7
Stem	40 years	1.711	2.701	0.929	0.024	90.0
	20 years	1.756	2.434	0.98	0.021	92.1
Total branch	40 years	-1.938	4.645	0.978	0.020	90.9
	20 years	1.104	2.566	0.949	0.062	84.5
New branch	40 years	-2.569	3.936	0.964	0.024	91.3
	20 years	-0.682	2.884	0.659	0.754	40.4
Older branch	40 years	-1.961	4.655	0.980	0.018	92.7
	20 years	1.105	2.562	0.957	0.051	85.9
Total foliage	40 years	-2.801	4.147	0.965	0.026	90.1
	20 years	1.437	1.714	0.867	0.079	82.5
Below ground	40 years	0.480	3.116	0.918	0.032	89.1
	20 years	0.272	2.674	0.948	0.069	82.8
Root	40 years	-3.008	4.335	0.991	0.007	95.1
	20 years	-1.641	2.322	0.981	0.081	91.9

R^2 , RSS , AA are respectively 0.88, 0.007, 84.1%. The R_{sbb} , R_{sb} and R_s harvesting regimes are widely used (Merino, 2005). Some organs (e.g., stalks, xylem of the trunk, leaves and bark) are directly related to the export of nutrients caused by harvesting, while other organs (e.g., roots) are unrelated to the export of nutrients caused by harvesting. Regardless of how old the larch trees were, among the organs that are directly related to the export of nutrients caused by harvesting, those organs (e.g., leaves, bark, new branches) [41] that exhibited a relatively high proportion of living tissue (e.g., parenchyma cells) showed relatively high nutrient contents (Fig. 1), whereas those (e.g., xylem) [41] that displayed a relatively low proportion of living tissue exhibited relatively low nutrient element contents. These results are similar to those of previous studies on other tree species [42], [43], [27].

The organs of the 20-year-old larch trees basically showed the same nutrient distribution patterns as in the 40-year-old larch trees, except for some individual indexes (e.g., N content, underground P content). There was no significant difference in the nutrient contents of the organs between the larch trees with two different stand ages (Fig. 1). However, due to the difference in biomass partitioning, there was a significant difference in the cumulative amounts of nutrients in the organs between the larch trees with two different stand ages (Fig. 1).

As the growth time extends, the cumulative amount of xylem biomass in the trunk of a tree will gradually increase [42], [43], and the proportion of the biomass partitioned to the xylem of the trunk of the tree will also gradually increase. As a result of the increase in the proportion of the biomass partitioned to the xylem of the trunks of the 40-year-old larch trees, the cumulative amounts of nutrients in the 40-year-old larch trees were significantly higher than in the 20-year-old larch trees. Similar to the xylem of the trunk, there was also a difference in the cumulative amounts of nutrients between the leaves and old branches due to the difference in biomass partitioning. However, in contrast to the xylem, the cumulative amounts of nutrient elements in the leaves and old branches of the 20-year-old larch trees were significantly higher than in the 40-year-old larch trees (Fig. 1). While there was no significant difference in the cumulative amounts of nutrient elements in the new branches and bark between the larch trees with two different stand ages, the difference in the cumulative amounts of nutrient elements in other organs resulted in a significant difference between the 20-year-old larch trees and the 40-year-old larch trees in terms of the aboveground cumulative total amount of nutrients (corresponding to the R_{sbb} harvesting regime), the cumulative amounts of nutrient elements in the trunk (corresponding to

the Rsb harvesting regime) and the cumulative amounts of nutrient elements in the xylem of the trunk (corresponding to the Rs harvesting regime) (Fig. 1).




Fig. 1. Contents and cumulative amounts of nutrients in each organ of the larch trees.

A: Bark; B: New branch; C: Older branch; D: Woody tissues of Stem; E: Total foliage; F: Below ground; G: Root; H: Total; I: Above ground; J: Stem; K: Total branch; L: Total branch +Total foliage

a: N concentration; b: P concentration; c: K concentration; d: Ca concentration; e: Mg concentration

a': N accumulation; b': P accumulation; c': K accumulation; d': Ca accumulation; e': Mg accumulation

3.3. Nutrient Contents in Tree Biomass Components and Soil

Compared with the Rsbb₂₀, prolonging the harvesting cycle (Rsbb₄₀) or switching to an Rsb₂₀ dramatically reduced the annual mean amounts of nutrients exported. Compared with the Rsbb₂₀, these two measures reduced the total amount of nutrients exported by 50.53–51.21% (reducing the annual mean amount of N exported by 71.30–75.54%, the annual mean amount of P exported by 35.07–37.69%, the annual mean amount of K exported by 25.49–34.85%, the annual mean amount of Ca exported by 34.13–34.95% and the annual mean amount of Mg exported by 44.86–50.19%) (Table 3). These measures also resulted in a decrease in the ratio of the annual mean total amount of nutrients exported to the annual net cumulative total amount of nutrients from 2.72 to 1.35–

1.33. Furthermore, these measures caused significant decreases in the proportions of the annual mean amount of each nutrient exported within the corresponding soil nutrient contents of the 0–30 cm soil layer (the proportion of the annual mean total amount of nutrients exported within the total soil nutrient content of the 0–30 cm soil layer decreased from 0.07% to 0.03%; that of N exported within the N content of this layer decreased from 0.07% to 0.02%; that of P exported within the P content of this layer decreased from 0.14% to 0.08–0.09%; that of K exported within the K content of this layer decreased from 0.06% to 0.04–0.05%; that of Ca exported within the Ca content of this layer decreased from 0.04% to 0.03%; and that of Mg exported within the Mg content of this layer decreased from 0.40% to 0.22–0.19%) (Table 3).

Table 3. Annual export of different nutrient elements in approximately 20- and 40-year-old larch plantation forests under different harvesting regimes.

		Rsbb20 years (Base type)	Rsb 20 years	Rs 20 years	Rsbb 40 years	Rsb 40 years	Rs 40 years
N	Annual nutrient exports (kg/hm ²)	13.65	3.92	1.87	3.34	2.57	1.56
	Annual nutrient exports/Annual nutrient exports of Base type	100.00%	28.70%	13.73%	24.46%	18.79%	11.44%
	Annual nutrient exports/Soil nutrients concentration (0-10cm)	0.23%	0.07%	0.03%	0.06%	0.04%	0.03%
	Annual nutrient exports/Soil nutrients concentration (0-20cm)	0.11%	0.03%	0.02%	0.03%	0.02%	0.01%
	Annual nutrient exports/Soil nutrients concentration (0-30cm)	0.07%	0.02%	0.01%	0.02%	0.01%	0.01%
P	Annual nutrient exports (kg/hm ²)	5.35	3.33	2.60	3.47	3.17	2.72
	Annual nutrient exports/Annual nutrient exports of Base type	100.00%	62.31%	48.58%	64.93%	59.32%	50.84%
	Annual nutrient exports/Soil nutrients concentration (0-10cm)	0.54%	0.34%	0.26%	0.35%	0.32%	0.28%
	Annual nutrient exports/Soil nutrients concentration (0-20cm)	0.24%	0.15%	0.11%	0.15%	0.14%	0.12%
	Annual nutrient exports/Soil nutrients concentration (0-30cm)	0.14%	0.08%	0.07%	0.09%	0.08%	0.07%
K	Annual nutrient exports (kg/hm ²)	3.72	2.43	1.87	2.77	2.68	2.33
	Annual nutrient exports/Annual nutrient exports of Base type	100.00%	65.15%	50.32%	74.51%	71.86%	62.64%
	Annual nutrient exports/Soil nutrients concentration (0-10cm)	0.17%	0.11%	0.09%	0.13%	0.12%	0.11%
	Annual nutrient exports/Soil nutrients concentration (0-20cm)	0.08%	0.05%	0.04%	0.06%	0.06%	0.05%
	Annual nutrient exports/Soil nutrients concentration (0-30cm)	0.06%	0.04%	0.03%	0.05%	0.04%	0.04%
Ca	Annual nutrient exports (kg/hm ²)	9.17	6.04	5.08	5.96	5.69	5.07
	Annual nutrient exports/Annual nutrient exports of Base type	100.00%	65.87%	55.46%	65.05%	62.03%	55.32%
	Annual nutrient exports/Soil nutrients concentration (0-10cm)	0.19%	0.13%	0.11%	0.13%	0.12%	0.11%
	Annual nutrient exports/Soil nutrients concentration (0-20cm)	0.09%	0.06%	0.05%	0.06%	0.05%	0.05%
	Annual nutrient exports/Soil nutrients concentration (0-30cm)	0.04%	0.03%	0.02%	0.03%	0.03%	0.02%
Mg	Annual nutrient exports (kg/hm ²)	1.06	0.58	0.39	0.53	0.50	0.39
	Annual nutrient exports/Annual nutrient exports of Base type	100.00%	55.14%	36.82%	49.81%	47.06%	36.44%
	Annual nutrient exports/Soil nutrients concentration (0-10cm)	2.43%	1.34%	0.90%	1.21%	1.14%	0.89%
	Annual nutrient exports/Soil nutrients concentration (0-20cm)	0.76%	0.42%	0.28%	0.38%	0.36%	0.28%
	Annual nutrient exports/Soil nutrients concentration (0-30cm)	0.40%	0.22%	0.15%	0.20%	0.19%	0.15%
Total nutrient	Annual nutrient exports (kg/hm ²)	32.95	16.30	11.82	16.08	14.60	12.07
	Annual nutrient exports/Annual nutrient exports of Base type	100.00%	49.47%	35.87%	48.79%	44.30%	36.63%
	Annual nutrient exports/Soil nutrients concentration (0-10cm)	0.24%	0.12%	0.09%	0.12%	0.11%	0.09%
	Annual nutrient exports/Soil nutrients concentration (0-20cm)	0.11%	0.05%	0.04%	0.05%	0.05%	0.04%
	Annual nutrient exports/Soil nutrients concentration (0-30cm)	0.07%	0.03%	0.02%	0.03%	0.03%	0.02%
	Annual nutrient exports/Annual nutrient accumulation of whole system*	2.72	1.35	0.98	1.33	1.21	1.00
	Annual production of Woody tissues (kg/hm ²)	1041.01			1285.31		

* Net cumulative total amount of nutrients of Larch plantation ecosystem equals to the difference value of nutrient content of the Runoff input and output (12.1 kg/hm², from Ding et al (Ding et al., 1989))

Compared with the Rsbb₂₀, the Rsb₄₀ or the Rs₄₀ further reduced the total amount of nutrients exported by 55.70–63.37% (reducing the annual mean amount of N exported by 81.21–88.56%, the annual mean amount of P exported by 40.68–49.16%, the annual mean amount of K exported by 28.14–37.36%, the annual mean amount of Ca exported by

37.97–44.68 and the annual mean amount of Mg exported by 52.94–63.56%) and the ratio of the annual mean total amount of nutrients exported to the annual net cumulative total amount of nutrients to 1.21–1.00 (Table 3). In addition, the proportions of the annual mean amounts of nutrients exported in the corresponding soil nutrient contents in the 0–30 cm soil layer also

continued to decrease.

Compared with the Rs_{bb20} , among the other harvesting regimes, the Rs_{20} resulted in the lowest annual mean amounts of nutrients exported. Under the Rs_{20} , the annual mean total amount of nutrients exported was only 35.87% of that under the Rs_{bb20} (with the annual mean amount of N exported decreasing by 86.27%; that of P exported decreasing by 51.42%; that of K exported decreasing by 49.68%; that of Ca exported decreasing by 44.54%; and that of Mg exported decreasing by 63.18%) (Table 3). In addition, the ratio of the annual mean total amount of nutrients exported to the annual net cumulative total amount of nutrients also decreased to 0.98. The proportions of the annual mean amounts of nutrients exported within the corresponding soil nutrient contents in the 0–30 cm soil layer also decreased to the lowest level (the proportion of the annual mean total amount of nutrients exported within the total soil nutrient content of the 0–30 cm soil layer decreased from 0.07% to 0.02%; that of N exported within the N content of this layer decreased from 0.07% to 0.01%; that of P exported within the P content of this layer decreased from 0.14% to 0.07%; that of K exported within the K content of this layer decreased from 0.06% to 0.03%; that of Ca exported within the Ca content of this layer decreased from 0.04% to 0.02%; and that of Mg exported within the Mg content of this layer decreased from 0.40% to 0.15%) (Table 3).

Compared with the harvesting regime with a 20-year harvesting cycle, the annual mean growth of the xylem increased from 1,041.01 kg/hm² to 1,285.31 kg/hm² under the harvesting regime with a 40-year harvesting cycle (Table 3).

4. Discussion

4.1. Construction of an Allometric Growth Equation

Allometric growth equations are a widely used, reliable method for estimating the biomass, net primary productivity and carbon budgets of trees [44]. However, there is currently no uniform model for allometric growth equations for trees. The majority of researchers accept a power function of $M=aD^b$ [45], [46], [47]. The allometric growth equations established in the present study were similar to those established in previous studies [46], [48]. The power function of $M=aD^b$ could be used to establish relatively satisfactory allometric growth equations, except for the new branches of the 20-year-old trees, similar to the results of previous studies on the growth of plants of the genus *Larix* [43], [27]. New branches are relatively more significantly affected by environmental factors. Therefore, relatively high variability of new branches has been observed in various tree species [49], [50], [43]. The new branches of larch trees in stands with relatively young ages exhibit particularly high variability. Some parameters of the new branches of larch trees in stands with relatively young ages, such as the total number of branches, branch diameter and branch angle, are not significantly correlated with DBH [34], which might be the main reason that more factors were required to fit the new branches of the 20-year-old larch trees.

4.2. Proportion of Belowground Biomass of Plants

Previous studies have shown that the belowground biomass of a plant accounts for approximately 21% of its total biomass, while the aboveground living biomass of a plant accounts for approximately 70% of its total biomass [51], [52]. King et al. studied the aboveground and belowground biomass of red pine trees in stands with different ages using the whole harvesting method and found that the belowground biomass accounted for approximately 22.64% of the biomass of the whole tree [53]. Our research results indicated that the proportion of belowground biomass in the larch plantations in northeastern China was relatively low (the belowground biomass of the 40-year-old and 20-year-old larch trees accounted for 15.6% and 16.8% of the total biomass, respectively) (Fig. 1), which might be due to the following reasons:

(1) Trees can adapt to environmental changes through adjusting their aboveground and belowground biomass structure. The type, age and size of a plant as well as the external environment can alter the partitioning pattern of the biomass of each organ in a forest tree [54]; i.e., plants adapt to changes in the nutrient environment through adjusting the aboveground and belowground partitioning patterns and maintain and balance the aboveground–belowground relationship through their feedback mechanism [55], [56]. When there is a lack of soil nutrients and water, a plant partitions more C to its belowground parts, resulting in an increase in the ratio of the belowground biomass to the aboveground biomass [57]. Previous researchers have observed this adaptive response in a number of plants [58], [59], [60]. For example, a hybrid poplar tree was found to partition more C to its aboveground parts due to a relatively high N supply [61], while P deficiency and low P conditions resulted in an increase in the proportion of carbohydrates partitioned to the roots by seedlings [62], [63]. In the present study, the soil contained abundant nutrients, which might be one of the main causes of the relatively low belowground biomass.

(2) The proportion of fine roots in the biomass of a larch tree can reach approximately 10% [64]. However, it is relatively difficult to collect samples from the belowground root system when employing the harvesting method, and a considerable amount of fine roots will inevitably be lost during the sampling process [65], which might be another cause of the relatively low belowground biomass indicated by the results of the present study.

(3) Stand age is an important factor that affects biomass partitioning [64], [65], [66], [67]. The proportion of branches and leaves in the biomass of a tree decreases with increasing stand age, while the proportion of the trunk increases, as observed in the larch trees in the present study. There are multiple reasons why stand age affects the allometric growth equation. First, the development of the height and leaves of a tree is relatively important in the early stage of tree growth, but the growth of the stalks and coarse roots is more important in the late stage. As a result, trees in stands with different ages

exhibit different biomass partitioning strategies [68]. In addition, relatively large trees display a relatively small leaf area, resulting in a decrease in the proportion of the branches and leaves [69]. The results of the present study show that the biomass of the trunk continued to increase with increasing stand age, but there was no significant difference in the total biomass of branches and leaves between trees in stands with different ages. Under the impact of these factors, the increase in the proportion of the trunk with increasing stand age will become a general phenomenon [70], [71], [72], [73].

4.3. Nutrient Element Distribution Patterns

There is relatively high similarity between the nutrient distribution patterns in different tree species in different regions. Generally speaking, leaves, small branches, bark and fine roots exhibit relatively high nutrient contents, whereas stalks, old branches and root systems show relatively low nutrient contents [74], [75], [76]. However, the nutrient distribution pattern also varies to a certain extent between species. For example, the leaves of larch trees display relatively low nutrient contents.

The environmental temperature is an important factor that affects the nutrient contents of leaves. Relatively low temperatures have an inhibitory effect on the degradation and mineralization rate of litter, thereby reducing the contents of N and P that soils can absorb [77]. In addition, relatively low temperatures reduce the capacity of a plant to absorb nutrients through altering the physiological characteristics of the plant (e.g., the permeability and liquid-state viscosity of the plasma membrane) [78], [79], and N and P contents exhibit a decreasing trend with decreasing temperatures. Northeastern China, a region suitable for the growth of larch trees, is characterized by relatively low temperatures, and plants in this region adapt to relatively low temperatures through relatively low N contents at the leaf level [80], resulting in relatively low nutrient contents of the leaves of larch trees.

4.4. Effect of Harvesting on Soil Nutrients

The nutrients lost by a forest ecosystem can be replenished through internal and external processes (e.g., precipitation, biological fixation) [81], [4], [82]. However, in a plantation ecosystem, the export of nutrients caused by harvesting may exceed the natural import of nutrients, resulting in a decrease in soil nutrients [7], [8]. The mechanism by which harvesting results in the loss of soil nutrients in a plantation is very complicated [16], as are the specific ways in which soil nutrients are lost [83], [2], [84]. Changes in the material cycle and hydrothermal conditions are some of the possible mechanisms by which harvesting results in a loss of soil nutrients [85], [86], [87] [88]. However, the important role of harvesting in the loss of soil nutrients in a plantation has gradually been recognized [7], [8]. The export of nutrients and a high proportion of soil nutrients will affect the long-term accumulation of soil nutrients [13]. The results of the present study showed that in the temperate larch plantations of northeastern China, the proportions of the total amount of

nutrients exported due to harvesting within the cumulative total amount of soil nutrients and the net cumulative total amount of nutrients in the system were both high and that the export of nutrients caused by harvesting would result in a significant decline in soil fertility (Table 3). Similar to findings in other plantations, it was shown in the present study that prolonging the harvesting cycle and changing the harvesting regime are both important methods for reducing the annual mean amounts of nutrients exported and the loss of soil nutrients [89]. The difference in the annual mean amount of nutrients exported between different harvesting ages was mainly due to the difference in the annual mean amount of nutrients exported from the branches, leaves and bark. However, the impact of the annual mean amount of nutrients exported from the xylem was relatively insignificant. Compared with the trunk, the branches, leaves and bark display a relatively short growth time [41]. Therefore, there is a relatively small difference in the biomass of the branches, leaves and bark between trees in stands with different ages. As a result, there is no significant difference in the annual mean amounts of nutrients exported from the branches, leaves and bark between different harvesting ages. Other tree species in northeastern China also have this characteristic [43]. Among the examined harvesting regimes, the annual mean amount of nutrients exported was significantly higher than the net cumulative amount of nutrients in the system under both the Rsbb and Rsb harvesting regimes, regardless of whether the 20-year or 40-year harvesting cycle was selected. The annual mean amount of nutrients exported was the same as the net cumulative amount of nutrients in the system under the Rs₄₀ and was slightly lower than the net cumulative amount of nutrients in the system under the Rs₂₀. While the Rs harvesting regime has been widely used internationally [3], this harvesting regime is not stipulated in the operational rules for forest harvesting of China. The Rs harvesting regime is not used in the harvesting in the forest areas in northeastern China. However, the Rs harvesting regime can effectively reduce the amounts of nutrients exported from plantations.

5. Conclusions

In the larch plantation ecosystems of northeastern China, the harvesting regime is an important factor resulting in the export of nutrients, which in turn causes declining soil fertility. Prolonging the harvesting cycle and adopting an Rs harvesting regime are both important methods for reducing the export of nutrients. The Rs₂₀ resulted in the lowest amount of nutrients being exported. However, there was no significant difference in the amount of nutrients exported between the Rs₂₀ and the Rs₄₀, mainly because the annual mean amounts of P and K exported under the Rs₂₀ were lower than those under the Rs₄₀, and the annual mean amounts of N, Ca and Mg exported under the Rs₂₀ were higher than those under the Rs₄₀. The Rs₄₀ resulted in a higher yield of timber. A harvesting regime should be selected based on the consideration of multiple factors, such as the results of the present study, the goal of harvesting (e.g., final harvesting, thinning or sanitation

harvesting) and the purpose of harvesting (e.g., for timber, firewood or craft material).

Acknowledgements

This research was supported by grants from National Natural Science Foundation of China (31370494, 41575137, 31170421, 31070406), and the key projects of Heilongjiang Province Natural Science Foundation (ZD201406), the National Basic Research Priorities Program of the Ministry of Science and Technology of China (2014FY110600).

References

- [1] Raison R J, Woods P V, Jakobsen B F, et al. 1986. Soil temperatures during and following low-intensity prescribed burning in a Eucalyptus pauciflora forest. *Soil Research*, 24 (1): 33–47.
- [2] Evans, J. 1999b. Sustainability of plantation forestry: impact of species change and successive rotations of pine in the Usutu Forest, Swaziland. *S. Afr. For. J.* 184, 63–70.
- [3] Merino A, Balboa M A, Soalleiro R R, et al. 2005. Nutrient exports under different harvesting regimes in fast-growing forest plantations in southern Europe. *Forest Ecology and Management*, 207 (3): 325–339.
- [4] Fisher, R. F., Binkley, D., 2000. *Ecology and Management of Forest Soils*. Wiley, New York.
- [5] FAO. 2010. *Global forest Resources Assessment 2010-Main Report*. Rome: Food and Agricultural Organization, 2010; 1–340.
- [6] Dambrine, E., Vega, J. A., Taboada, T., Rodriguez, L., Fernandez, C., Macias, F., Gras, J. M. 2000. Bilans d'éléments minéraux dans de petits bassins versants forestiers de Galice (NW Espagne). *Annals of forest science*, 57 (1): 23–38.
- [7] Johnson, D. W., Todd, D. E., 1998. Harvesting effects on long-term changes in the nutrient pool of mixed oak forest. *Soil Sci. Soc. Am. J.* 62, 1725–1735.
- [8] Olsson, B. A., Lundkvist, H., Staaf, H., 2000. Nutrient status in needles of Norway spruce and Scots pine following harvesting of logging residues. *Plant Soil* 23, 161–173.
- [9] Gómez, M. X., Calvo de Anta, R., 2001. Ciclo de agua y elementos en suelos forestales (*Pinus radiata*) de Galicia. III Congreso Forestal Español, mesa 2. pp. 567–572.
- [10] Miller, H. G., 1984. Dynamics of nutrient cycling in plantation ecosystems. In: Bowen, G. D., Nambiar, E. K. S. (Eds.), *Nutrition of Plantation Forests*. Academic Press, London, pp. 53–78.
- [11] Wisconsin Department of Natural Resources. 2008. *Wisconsin's Forestland Woody Biomass Harvesting Guidelines*. Madison, WI).
- [12] Walker T, Cardellicchio P, Colnes A, et al. 2010. Biomass sustainability and carbon policy study. Manomet Center for Conservation Sciences.
- [13] Fassbender, H., Bornemisza, E., 1987. *Química de Suelos con Énfasis en Suelos de América Latina*. Instituto Interamericano de Cooperación para la Agricultura (IICA), San José, p. 420.
- [14] Vitousek, P., 1982. Nutrient cycling and nutrient use efficiency. *Am. Naturalist*, 119, 553–572.
- [15] Vitousek, P., 1984. Litter, nutrient cycling, and nutrient limitation in tropical forests. *Ecology*, 65, 285–298.
- [16] Wang, D., Bormann, F. H., Lugo, A. E., Bowden, R. D., 1991. Comparison of nutrient-use efficiency and biomass production in five tropical tree taxa. *Forest Ecology and Management*, 46, 1–21.
- [17] Kumar, B. M., Geoga, S. J., Jamaludheen, V., Suresh, T. K., 1998. Comparison of biomass production, tree allometry and nutrient use efficiency of multipurpose trees grown in woodlot and silvopastoral experiments in Kerala, India. *Forest Ecology and Management*, 112, 145–163.
- [18] Smith, K., Gholz, H. L., Oliveira, F. A., 1998. Litter fall and nitrogen-use efficiency of plantations and primary forest in the eastern Brazilian Amazon. *Forest Ecology and Management*, 109, 209–220.
- [19] Callaway R M, Walker L R. 1997. Competition and facilitation: a synthetic approach to interactions in plant communities. *Ecology*, 78 (7): 1958–1965.
- [20] Fölster, H., Khanna, P. K., 1997. Dynamics of nutrient supply in plantations soils. In: Nambiar, E. K. S., Brown, A. G. (Eds.), *Management of Soil Nutrient and Water in Tropical Plantation Forests*. Australian Centre for International Agricultural Research, Canberra, pp. 339–378.
- [21] Chen Xi-quan. 1986. The effect of community structure of artificial larch wood on physico-chemical properties of Albic soils. *Journal of northeast forestry university*, 4: 113–116.
- [22] Chen Zhiguo Wang Shuchen Che Zhiwen. 1991. Study of both *Larix gmelini* × *Fraxinus mandshurica* and *Larix gmelini* × *juglans mandshurica* mixed stand. *Journal of Northeast Forestry University*, 1: 100–105.
- [23] Wang Hongjun, Gong Fang, Zheng Baoren, Chen Dexiang. 1997. Physical and Chemical Properties of the Soil for Larch Plantations. *Journal of northeast forestry university* 25: 75–79.
- [24] Yang K, Zhu J-J, Zhang J-X, et al. 2009. Seasonal dynamics of soil microbial biomass C and N in two larch plantation forests with different ages in Northeastern China. *Acta Ecologica Sinica*, 29 (10): 5500–5507.
- [25] Yang K, Zhu JJ, Zhang M, et al. 2010. Soil microbial biomass carbon and nitrogen in forest ecosystems of Northeast China: A comparison between natural secondary forest and larch plantation. *Journal of Plant Ecology*, 3: 175–182.
- [26] Yang K, Zhu JJ, Yan QL, et al. 2013. Soil enzyme activities as potential indicators of soluble organic nitrogen pools in forest ecosystems of Northeast China. *Annals of Forest Science*, 69: 795–803.
- [27] Yan Tao, Zhu Jiao-jun, Yang Kai, Yu Li-zhong. 2014. Chinese Aboveground biomass and nutrient distribution patterns of larch plantation in a montane region of eastern Liaoning Province, China. *Journal of Applied Ecology*, 10: 2772–2778.
- [28] Mei L, Zhang Z-W, Gu J-C, et al. 2009. Carbon and nitrogen storages and allocation in tree layers of *Fraxinus mandshurica* and *Larix gmelinii* plantations. *Chinese Journal of Applied Ecology*, 20 (8): 1791–1796.

- [29] Mrino A, Balboa MA, Soalleiro RR, et al. 2005. Nutrient exports under different harvesting regimes in fast-growing forest plantations in southern Europe. *Forest Ecology and Management*, 207: 325–339.
- [30] Raulund-Rasmussen K, Stupak I, Clarke N, et al. 2008. Effects of very intensive forest biomass harvesting on short and long term site productivity // Rser D, ed. *Sustainable Use of Forest Biomass for Energy: A Synthesis with Focus on the Baltic and Nordic Region*. Amsterdam, the Netherlands: Springer: 56–67.
- [31] Thiffault E, Hannam KD, Paré D, et al. 2011. Effects of forest biomass harvesting on soil productivity in boreal and temperate forests: A review. *Environmental Reviews*, 19: 278–309.
- [32] Feng Z-W, Wang X-K, Wu G. 1999. *Biomass and Productivity of Forest Ecosystems in China*. Beijing: Science Press.
- [33] Li Y-K. 1989. *Soil Agricultural and Chemical Routine Analysis Method*. Beijing: Science Press, 1989.
- [34] Zhang H, Guan DS, Song MW. 2012. Biomass and carbon storage of Eucalyptus and Acacia plantations in the Pearl River Delta, South China. *Forest Ecology and Management*, 277: 90–97.
- [35] Kelly W, Brittney R, James B. 2013. Effects of timber harvest intensity on macronutrient cycling in oak-dominated stands on sandy soils of northwest Wisconsin. *Forest Ecology and Management*, 291: 1–12.
- [36] Marcin P, Barthlomej W, Nicholas H. 2013. Scots pine needles macronutrient (N, P, K, Ca, Mg and S) supply at different reclaimed mine soil substrates: As an indicator of the stability of developed forest ecosystems. *Environmental Monitoring and Assessment*, 185: 7445–7457.
- [37] Feldpausch T R, Rondon M A, Fernandes E C M, et al. 2004. Carbon and nutrient accumulation in secondary forests regenerating on pastures in central Amazonia. *Ecological Applications*, 14 (sp4): 164–176.
- [38] Peri P L, Gargaglione V, Pastur G M. 2008. Above- and belowground nutrients storage and biomass accumulation in marginal Nothofagus antarctica forests in Southern Patagonia. *Forest Ecology and Management*, 255: 2502–2511.
- [39] Hou L, Lei R-D. 2009. Carbon dioxide sequestration of main shrub species in a natural secondary Pinus tabulaeform forest at the Huoditang forest zone in the Qinling Mountains. *Acta Ecologica Sinica*, 29 (11): 6077–6084.
- [40] Li Y-Q, Zheng S-W, Su Y-M, et al. 2009. Growth and allometric biomass equations of introduced Agave americana L. in upper reaches of Minjiang River. *Acta Ecologica Sinica*, 29 (9): 4820–4826.
- [41] Esau K. 1977. *The anatomy of seed plants*. Ed. Wiley & Sons, New York.
- [42] West G B, Brown J H, Enquist B J. 2001. A general model for ontogenetic growth. *Nature*, 413 (6856): 628–631.
- [43] Wang C. 2006. Biomass allometric equations for 10 co-occurring tree species in Chinese temperate forests. *Forest Ecology and Management*, 222 (1): 9–16.
- [44] Gower, S. T., Kucharik, C. J., Norman, J. M., 1999. Direct and indirect estimation of leaf area index, F (APAR), and net primary production of terrestrial ecosystems. *Remote Sens. Environ.* 70, 29–51.
- [45] Niklas, K. J. 1994. *Plant Allometry: The Scaling of Form and Process*. University of Chicago Press, Chicago, Illinois.
- [46] Ter-Mikaelian, M. T., Korzukhin, M. D., 1997. Biomass equations for sixty-five North American tree species. *Forest Ecology and Management*, 97, 1–24.
- [47] Ketterings, Q. M., Coe, R., Van Noordwijk, M., Ambagau, Y., Palm, C. A. 2001. Reducing uncertainty in the use of allometric biomass equations for predicting above-ground tree biomass in mixed secondary forests. *Forest Ecology and Management*, 146, 199–209.
- [48] Bond-Lamberty, B., Wang, C., Gower, S. T. 2002. Aboveground and below-ground biomass and sapwood area allometric equations for six boreal tree species of northern Manitoba. *Can. J. For. Res.* 32, 1441–1450.
- [49] Crow, T. R., and Schlaegel, B. E. 1988. A guide to using regression equations for estimating tree biomass. *North. J. Appl. For.* 5: 15–22.
- [50] Buech, R. R., and Rugg, D. J. 1989. Biomass relations of shrub components and their generality. *Forest Ecology and Management*, 26: 257–264.
- [51] Houghton RA, Lawrence KT, Hackler JL, Brown S. 2001. The spatial distribution of forest biomass in the Brazilian Amazon: a comparison of estimates. *Global Change Biology*, 7 (7): 731–746.
- [52] Saatchi SS, Houghton RA, Dos Santos Alvala RC, Soares JV, Yu Y. 2007. Distribution of aboveground live biomass in the Amazon basin. *Global Change Biology*, 13 (4): 816–837.
- [53] King JS, Giardina CP, Pregitzer KS, Friend AL. 2007. Biomass partitioning in red pine (*Pinus resinosa*) along a chronosequence in the Upper Peninsula of Michigan. *Can J For Res*, 37 (1): 93–102.
- [54] Son Y, Hwang J W, Kim Z S, et al. 2001. Allometry and biomass of Korean pine (*Pinus koraiensis*) in central Korea. *Bioresource Technology*, 78 (3): 251–255.
- [55] Ericsson T, Ingestad T. 1988. Nutrition and growth of birch seedlings at varied relative phosphorus addition rates. *Physiologia Plantarum*, 72 (2): 227–235.
- [56] Ingestad T, Agren G I. 1991. The Influence of Plant Nutrition on Biomass Allocation. *Ecological Applications*, 1 (2): 168–174.
- [57] Fabião, A., Madeira, M., Steen, E., Kätterer, T., Ribeiro, C., & Araújo, C. 1994. Development of root biomass in an Eucalyptus globulus plantation under different water and nutrient regimes. *Plant & Soil*, 168-169 (1), 215–223.
- [58] Larigauderie A, Reynolds J F, Strain B R. 1994. Root response to CO₂ enrichment and nitrogen supply in loblolly pine. *Plant & Soil*, 165 (1): 21–32.
- [59] Mackiedawson L A, Millard P, Proe M F. 1995. The Effect Of Nitrogen Supply On Root Growth And Development In Sycamore And Sitka Spruce Trees. *Forestry*, 68 (2): 107–114.
- [60] Proe M F, Millard P. 1995. Effect of P supply upon seasonal growth and internal cycling of P in Sitka spruce (*Picea sitchensis* (Bong) carr) seedlings. *Plant & Soil*, 168-169 (1): 313–317.
- [61] Ibrahim L, Proe M F, Ad. C. 1998. Interactive effects of nitrogen and water availabilities on gas exchange and whole-plant carbon allocation in poplar. *Tree Physiology*, 18 (7): 481–487.

- [62] Topa M A, Jm. C. 1992. Carbon and phosphorus partitioning in *Pinus serotina* seedlings growing under hypoxic and low-phosphorus conditions. *Tree Physiology*, 10 (2): 195–207.
- [63] Proe M F, Millard P. 1995. Effect of N supply upon the seasonal partitioning of N and P uptake in young Sitka spruce (*Picea sitchensis*). *Canadian journal of forest research*, 25 (10): 1704–1709.
- [64] Rapp M, Santa Regina I, Rico M, et al. 1999. Biomass, nutrient content, litterfall and nutrient return to the soil in Mediterranean oak forests. *Forest Ecology and Management*, 119 (1): 39–49.
- [65] Yowhan Son, Jae Woo, H wang, Zin Suh Kim, Woo Kyun Lee, Jong Sung Kim. 2001. Allometry and biomass of Korean pine in central Korea. *Bioresource technology*, 2001 (78): 251–255.
- [66] Xiao CW, Ceulemans R. 2004. Allometric relationships for below- and aboveground biomass of young Scots pines. *Forest Ecology and Management*, 203 (1/3): 177–186.
- [67] Castilho CV, Magnusson WE, Araujo RN, Luizao RCC, Luizao FJ, Lima AP, Higuchi N. 2006. Variation in aboveground tree live biomass in a central Amazonian Forest: Effects of soil and topography. *Forest Ecology and Management*, 234: 85–96.
- [68] Peichl M, Arain M A. 2007. Allometry and partitioning of above- and belowground tree biomass in an age-sequence of white pine forests. *Forest Ecology and Management*, 253: 68–80.
- [69] Magnani F, Mencuccini M, Grace J. 2000. Age-related decline in stand productivity: the role of structural acclimation under hydraulic constraints. *Plant Cell Environ* 23: 251–263.
- [70] Satoo, T., Madgwick, H. A. I., 1982. *Forest Biomass*. M. Nijhoff/Dr. W. Junk Publishers, Boston, 152 pp.
- [71] Helmisaari, H.-S., Makkonen, K., Kellomaki, S., Valtonen, E., Malkonen, E., 2002. Below- and above-ground biomass, production and nitrogen use in Scots pine stands in eastern Finland. *Forest Ecology and Management*, 165, 317–326.
- [72] Montagu KD, Duttmer K, Barton CVM, Cowie AL. 2005. Developing general allometric relationships for regional estimates of carbon sequestration—an example using *Eucalyptus pilularis* from seven contrasting sites. *Forest Ecology and Management*, 204: 113–127.
- [73] Lee YJ, Lee MH, Lee KH, Son YM, Seo JH, Park IH, Son Y. 2006. Effects of stand age classes on biomass expansion factors and stem densities in *Chamaecyparis obtusa* plantations. *J Korean For Soc* 95: 50–54 (in Korean).
- [74] Caldentey, J., Schmidt, H., Bowen, H. 1993. Acumulación de biomasa y nutrientes en rodales de Lenga (*Nothofagus pumilio*) en Magallanes, Chile. In: *Proceedings of the VII Jornadas Técnicas Forestales: Ecosistemas Forestales Nativos—Uso, Manejo y Conservación*, El Dorado, Argentina, pp. 22–36.
- [75] Hart, P. B. S., Clinton, P. W., Allen, R. B., Nordmeyer, A. H., Evans, G. 2003. Biomass and macro- nutrients (above-and below-ground) in a New Zealand beech (*Nothofagus*) forest ecosystem: implications for storage and sustain-able forest management. *Forest Ecology and Management*, 174, 281–294.
- [76] Peri P L, Gargaglione V, Pastur G M. 2006. Dynamics of above-and below-ground biomass and nutrient accumulation in an age sequence of *Nothofagus antarctica* forest of Southern Patagonia. *Forest Ecology and Management*, 233 (1): 85–99.
- [77] Kirschbaum, M. U. F. 1995. The temperature dependence of soil organic matter decomposition and the effect of global warming on soil organic carbon storage. *Soil Biology and Biochemistry*, 27: 753–760.
- [78] Hobbie, S. E., Schimel, J. R., Trumbore, S. E. & Randerson, J. R. 2000. A mechanistic understanding of carbon storage and turnover in high-latitude soils. *Global Change Biology*, 6: 196–210.
- [79] Hobbie, S. E., Nadelhoffer, K. J. & Hogberg, R. 2002. A synthesis: the role of nutrients as constraints on carbon balances in boreal and arctic regions. *Plant and Soil*, 242: 163–170.
- [80] Han W, Fang J, Guo D, et al. 2005. Leaf nitrogen and phosphorus stoichiometry across 753 terrestrial plant species in China. *New Phytologist*, 168 (2): 377–385.
- [81] Verry E S, Timmons D R. 1977. Precipitation nutrients in the open and under two forests in Minnesota. *Canadian Journal of Forest Research*, 7 (1): 112–119.
- [82] Sparks J P. 2009. Ecological ramifications of the direct foliar uptake of nitrogen [J]. *Oecologia*, 159 (1): 1–13.
- [83] Woods, R. V. 1990. Second rotation decline in *P. radiata* plantations in South Australia has been corrected. *Water Air Soil Poll.* 54, 607–619.
- [84] McNabb, K. L., Wadouski, L. H., 1999. Multiple rotation yields for intensively managed plantations in the Amazon basin. *New For.*, 18 (1), 5–15.
- [85] Mroz GD, Jurgensen MF and Frederick DJ. 1985. Soil nutrient changes following whole tree harvesting on tree northern hardwood sites. *Soil Soc. Sci*, 49: 1552–1557.
- [86] Gessel S P, Tweruer J. 1990. Potential nutritional problems in main tenancy of long term southern hemisphere plant at ion productivity. *IUFRO 19th World Congress Proceedings*, Canada.
- [87] Dahlgren RA, Driscoll CT. 1994. The effects of whole tree clear cutting on soil processes at the Hubbard Brook experimental forest. *Plant Soil*, 158: 239–262.
- [88] Mohammad AH, Ahmad J and Hamid RK. 1997. Deforest at ion effects on soil physical and chemical properties, Lordegan, Ran. *Plant and Soil*, 190: 301–308.
- [89] Ma X, Heal K V, Liu A, et al. 2007. Nutrient cycling and distribution in different-aged plantations of Chinese fir in southern China. *Forest Ecology and Management*, 243 (1): 61–74.